

Considerations of Environmentally Relevant Test Conditions for Improved Evaluation of Ecological Hazards of Engineered Nanomaterials

Citation for published version:

Holden, PA, Gardea-Torresdey, JL, Klaessig, F, Turco, RF, Mortimer, M, Hund-Rinke, K, Cohen Hubal, EA, Avery, D, Barceló, D, Behra, R, Cohen, Y, Deydier-Stephan, L, Ferguson, PL, Fernandes, TF, Herr Harthorn, B, Henderson, WM, Hoke, RA, Hristozov, D, Johnston, JM, Kane, AB, Kapustka, L, Keller, AA, Lenihan, HS, Lovell, W, Murphy, CJ, Nisbet, RM, Petersen, EJ, Salinas, ER, Scheringer, M, Sharma, M, Speed, DE, Sultan, Y, Westerhoff, P, White, JC, Wiesner, MR, Wong, EM, Xing, B, Steele Horan, M, Godwin, HA & Nel, AE 2016, 'Considerations of Environmentally Relevant Test Conditions for Improved Evaluation of Ecological Hazards of Engineered Nanomaterials', *Environmental Science and Technology*, vol. 50, no. 12, pp. 6124-6145. <https://doi.org/10.1021/acs.est.6b00608>

Digital Object Identifier (DOI):

[10.1021/acs.est.6b00608](https://doi.org/10.1021/acs.est.6b00608)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Peer reviewed version

Published In:

Environmental Science and Technology

Publisher Rights Statement:

This document is the unedited author's version of a Submitted Work that was subsequently accepted for publication in Environmental Science and Technology, © American Chemical Society after peer review. To access the final edited and published work, see <http://pubs.acs.org/doi/abs/10.1021/acs.est.6b00608>

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

24 Development, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina
 25 27711, United States; ⁹Department of Environmental Chemistry, Institute of Environmental
 26 Assessment and Water Research (IDAEA-CSIC), Barcelona 08034, Spain; ¹⁰Institut Català de
 27 Recerca de l'Aigua (ICRA), Parc Científic i Tecnològic de la Universitat de Girona, Girona
 28 17003, Spain; ¹¹Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600
 29 Dübendorf, Switzerland; ¹²Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich,
 30 8092 Zürich, Switzerland; ¹³California NanoSystems Institute, ¹⁴Chemical and Biomolecular
 31 Engineering Department, University of California Los Angeles, California 90095, United States;
 32 ¹⁵European Chemicals Agency (ECHA), Helsinki 00210, Finland; ¹⁶Department of Civil &
 33 Environmental Engineering, ¹⁷Center for the Environmental Implications of NanoTechnology
 34 (CEINT), Duke University, Durham, North Carolina 27708, United States; ¹⁸Heriot-Watt
 35 University, Edinburgh EH14 4AS, United Kingdom; ¹⁹Center for Nanotechnology in Society,
 36 ²⁰Department of Anthropology, University of California, Santa Barbara, California 93106;
 37 ²¹Office of Research and Development, National Exposure Research Laboratory, U.S.
 38 Environmental Protection Agency, Athens, Georgia 30605, United States; ²²E.I. du Pont de
 39 Nemours and Company, Newark, Delaware 19711, United States; ²³Department of
 40 Environmental Sciences, Informatics and Statistics, University Ca' Foscari Venice, Venice
 41 30123, Italy; ²⁴Department of Pathology and Laboratory Medicine, Brown University,
 42 Providence, Rhode Island 02912, United States; ²⁵LK Consultancy, Turner Valley, Alberta T0L
 43 2A0, Canada; ²⁶Vive Crop Protection Inc, Toronto, Ontario M5G 1L6, Canada; ²⁷Department of
 44 Chemistry, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, United States;
 45 ²⁸Department of Ecology, Evolution and Marine Biology, University of California, Santa
 46 Barbara, California 93106, United States; ²⁹Biosystems and Biomaterials Division, Material

47 Measurement Laboratory, National Institute of Standards and Technology, Gaithersburg,
48 Maryland 20899, United States; ³⁰BASF SE, Experimental Toxicology and Ecology,
49 Ludwigshafen, D-67056, Germany; ³¹Institute for Chemical and Bioengineering, ETH Zürich,
50 8093 Zürich, Switzerland; ³²PETA International Science Consortium, Ltd., London N1 9RL,
51 England, United Kingdom; ³³Globalfoundries, Corporate EHS, Hopewell Junction, New York
52 12533, United States; ³⁴Environment Canada, Gatineau, Quebec J8X 4C8, Canada; ³⁵School of
53 Sustainable Engineering and the Built Environment, Arizona State University, Tempe, Arizona
54 85287, United States; ³⁶Department of Analytical Chemistry, The Connecticut Agricultural
55 Experiment Station, New Haven, Connecticut 06504, United States; ³⁷Office of Pollution
56 Prevention and Toxics, U.S. Environmental Protection Agency, Washington, D.C. 20460,
57 United States; ³⁸Stockbridge School of Agriculture, University of Massachusetts, Amherst,
58 Massachusetts 01003, United States; ³⁹Department of Environmental Health Sciences, Fielding
59 School of Public Health, ⁴⁰Institute of the Environment and Sustainability, ⁴¹Division of
60 NanoMedicine, Department of Medicine, University of California, Los Angeles, California
61 90095, United States

62

63 **Corresponding Author**

64 E-mail: holden@bren.ucsb.edu; tel: 805-893-3195; fax: 805-893-7612

65

66 **Author E-Mail Address**

67 Patricia A. Holden: holden@bren.ucsb.edu

68 Jorge F. Gardea-Torresdey: gardea@utep.edu

69 Frederick Klaessig: fred.klaessig@verizon.net

70 Ronald F. Turco: rturco@purdue.edu

71 Monika Mortimer: mmortimer@bren.ucsb.edu
72 Kerstin Hund-Rinke: kerstin.hund-rinke@ime.fraunhofer.de
73 Elaine A. Cohen Hubal: Hubal.elaine@Epa.gov
74 David Avery: avery@ucla.edu
75 Damià Barceló: dbcqam@cid.csic.es
76 Renata Behra: renata.behra@eawag.ch
77 Yoram Cohen: yoram@ucla.edu
78 Laurence Deydier-Stephan: laurence.deydier@echa.europa.eu
79 Patrick Lee Ferguson: lee.ferguson@duke.edu
80 Teresa Fernandes: t.fernandes@hw.ac.uk
81 Barbara Herr Harthorn: harthorn@cns.ucsb.edu
82 William Matthew Henderson: henderson.matt@epa.gov
83 Robert A. Hoke: robert.a.hoke@dupont.com
84 Danail Hristozov: danail.hristozov@unive.it
85 John M. Johnston: johnston.johnm@epa.gov
86 Agnes B. Kane: agnes_kane@brown.edu
87 Larry Kapustka: kapustka@xplornet.com
88 Arturo A. Keller: keller@bren.ucsb.edu
89 Hunter S. Lenihan: Lenihan@bren.ucsb.edu
90 Wess Lovell: wlovell@vivecrop.com
91 Catherine J. Murphy: murphycj@illinois.edu
92 Roger N. Nisbet: nisbet@lifesci.ucsb.edu
93 Elijah J. Petersen: elijah.petersen@nist.gov

94 Edward R. Salinas: Edward.salinas@basf.com
95 Martin Scheringer: scheringer@chem.ethz.ch
96 Monita Sharma: monitas@piscld.org.uk
97 David E. Speed: david.speed@globalfoundries.com
98 Yasir Sultan: yasir.sultan@canada.ca
99 Paul Westerhoff: p.westerhoff@asu.edu
100 Jason C. White: Jason.White@ct.gov
101 Mark R. Wiesner: wiesner@duke.edu
102 Eva M. Wong: wong.eva@epa.gov
103 Baoshan Xing: bx@umass.edu
104 Meghan Steele Horan: meghan.horan@ucla.edu
105 Hilary A. Godwin: hgodwin@ucla.edu
106 André E. Nel: anel@mednet.ucla.edu
107
108
109
110

111 **Abstract**

112 Engineered nanomaterials (ENMs) are increasingly entering the environment with
113 uncertain consequences including potential ecological effects. Various research
114 communities view differently whether ecotoxicological testing of ENMs should be
115 conducted using environmentally relevant concentrations—where observing outcomes is
116 difficult—versus higher ENM doses, where responses are observable. What exposure
117 conditions are typically used in assessing ENM hazards to populations? What conditions
118 are used to test ecosystem-scale hazards? What is known regarding actual ENMs in the
119 environment, via measurements or modeling simulations? How should exposure
120 conditions, ENM transformation, dose, and body burden be used in interpreting biological
121 and computational findings for assessing risks? These questions were addressed in the
122 context of this critical review. As a result, three main recommendations emerged. First,
123 researchers should improve ecotoxicology of ENMs by choosing test endpoints, duration,
124 and study conditions—including ENM test concentrations—that align with realistic
125 exposure scenarios. Second, testing should proceed via tiers with iterative feedback that
126 informs experiments at other levels of biological organization. Finally, environmental
127 realism in ENM hazard assessments should involve greater coordination among ENM
128 quantitative analysts, exposure modelers, and ecotoxicologists, across government,
129 industry, and academia.

Introduction

As nanotechnology (the synthesis, manipulation, and measurement of matter at 10^{-9} meter scales)¹ rapidly evolves,² engineered nanomaterials (ENMs) are entering air, waters, soils,³ and sediments where they could adversely affect organisms to ecosystems.⁴⁻⁶ Actual environmental impacts of ENMs have not been documented, and there are uncertainties about the potential for, and how to evaluate, impacts.⁷

ENM ecotoxicology elucidates hazards and their mechanisms.⁸ The scope overlaps with conventional ecotoxicology, although ENMs are particulate and diverse,⁹ with varying cores, native or acquired surface chemistries,¹⁰ conditional agglomeration or dissolution,¹¹ and size- plus composition-dependent electronic properties,¹² affecting their reactivity and biological interactions.¹³

Focusing ENM ecotoxicology invokes exposure scenarios^{14, 15} relevant to ENM production,¹⁶⁻¹⁹ use,²⁰⁻²⁴ disposal,^{25, 26} and product release (Fig. 1).^{14, 27-31} Scenarios consider environmental fate and transport,^{15, 32-39} bioavailability,^{40, 41} and ENM uptake into ecological receptors.⁴²⁻⁴⁶ Scenarios also specify processes^{47, 48} and ecosystem services⁴⁹ such as food production.^{50, 51}

In conventional chemical toxicology, observed and perceived exposures often diverge.⁵² In ENM ecotoxicology, water is emphasized, while soil and sediment impacts have received less attention.⁵³ Where do exposures occur? What ENM forms⁵⁴ and quantities are involved? Which ecological receptors are affected?⁵⁵ Which local exposure conditions prevail? As with conventional chemical risk assessment, such questions unite hazard and exposure assessments.^{15, 56}

Standardized test regimens do not derive from scenarios, since ENM test conditions are predefined (e.g., aqueous chemistry, temperature, pH, experiment duration, among others) for standardized endpoints.⁵⁷ Ideally, ENM hazards are studied at realistic exposures for ecologically relevant receptors. An example would be studying real soils under controllable yet realistic conditions, i.e. in greenhouses⁵⁸ or lysimeters.⁵⁹ However, requiring absolute realism in all ENM ecotoxicology would pose scientific challenges associated with: measuring ENMs analytically in environmental media; measuring toxicity across a representative range of environmental conditions; characterizing environmental ENM forms and their transformations so that toxicity is measured for representative materials; ENMs altering physical or chemical exposure media conditions; and few efficient approaches for estimating hazards and exposures necessary to evaluate risks before ENM products develop. Another challenge is internal to the scientific community: multiple dissimilar working definitions of environmental relevance intruding on scholarship, including peer review.⁵³ Environmental relevance remains undefined, leading to categorization of research around a few selected concepts.⁶⁰

Previously, over 600 published studies were examined to compare modeled or measured environmental concentrations of ENMs versus concentrations administered in ENM ecotoxicity assessments.⁵³ The study found nominal concentration disparities, but also infrequent testing at low ENM concentrations. The study noted uncertainties in ENM exposure modeling, and that other toxicity testing conditions beyond ENM concentration—including aqueous chemistry, biological receptor, system complexity, and ENM form—relate to real-world conditions. However, the study did not establish what constitutes environmental relevance in the ecotoxicology of ENMs.⁵³

Issues regarding environmental detection, uncertain concentrations, and unknown toxicity are not unique to ENMs; they are also raised for other emerging contaminants.⁶¹ However, because the ENM industry is rapidly evolving and scientists seek to assist in advancing environmentally safe nanotechnology, environmental relevance in ENM hazard assessment should be prioritized. To accomplish this, representatives from academia, industry, and government regulation working in ecotoxicology, exposure modeling, and social science (Table S1) addressed four questions within this critical review: (i) What exposure conditions are used in assessing ENM ecotoxicity potential for model organisms? (ii) What exposure and design considerations drive mesocosm experiments for assessments of ENM environmental hazards? (iii) What is the state of knowledge regarding ENM environmental exposure conditions, via measurements or modeling simulations? (iv) How should concepts such as exposure conditions, ENM transformation, dose, and body burden be used in interpreting biological and computational findings for assessing risks? The main objective was to provide context and guidance to the meaning of environmental relevance (singular or multiple meanings) in ENM environmental hazard assessment. This critical review addresses the four motivating questions and expands on detailed topics that emerged during the project (Supporting Information). For each question, there are findings and recommendations. These serve to crystallize what is meant by environmental relevance in ENM ecotoxicology, and further coalesce ENM environmental exposure and hazard assessment endeavors.

What exposure conditions are used in assessing ENM ecological hazard potential for model organisms?

Findings

Many systems, approaches, and conditions (including standardized testing with model media, organisms, and endpoints)⁶² have been used to assess ENM ecotoxicity.⁶³ The applicability and challenges of standardized testing protocols under aquatic conditions have been reviewed^{57, 64} and vetted in workshops.⁶² Also, the OECD reviewed and vetted its guidelines for testing ENMs⁶⁵ in an expert meeting.^{66, 67} Readers are referred to those reports for deliberations of ENM standardized testing.

Research studies include laboratory-specific low-^{68, 69} and high-throughput dose-response evaluations using select media with ENM compositional and receptor variants⁷⁰ for assessing uptake⁷¹⁻⁷³ and effect mechanisms.^{69, 74, 75} Investigations include mechanistic gene transcriptional,⁷⁶⁻⁷⁹ DNA damage,⁸⁰ metabolomics profiling,⁸¹ and transgenerational^{82, 83} experiments. Bottle-scale microcosms partially simulate limited levels of environmental complexity,⁸⁴ e.g. in using natural soils or sediments with⁵⁸ or without^{85, 86} plants and associated rhizosphere influences,⁸⁷ or using seawater⁸⁸ or marine sediments^{89, 90} and associated receptors based on expected ENM compartmentalization.⁹¹ Non-standard microcosms assess ecological endpoints, e.g. microbial community composition^{85, 92} and function⁹² related to C and N cycling.^{92, 93} Environmental factors are examined, including how ENMs interactively affect soil water availability⁹⁴ and soil bacterial communities.⁹⁵ Single-species experiments that assess ENM bio-association^{96, 97} or bioaccumulation^{71, 98} precede and motivate using microcosms for assessing dual species trophic transfer⁹⁸⁻¹⁰¹ and potential biomagnification.¹⁰²

Pristine ENMs, including those with surface functionalization, capping agents, or adsorbed species or coatings,¹⁰³ are the most frequently assessed, although released and

transformed versions are increasingly studied.^{28, 30, 104} Results for textiles, paints, and nanocomposites¹⁰⁵ suggest that released particles significantly transform and age, and exhibit different environmental behavior and effects compared to pristine ENMs.^{106, 107} Assessing changes in form and associated behavior or activity across the material's life cycle are uniquely challenging.

Nominal test exposure concentrations vary widely⁵³ (Fig. S1) and are sometimes related to scenarios such as repeated applications,¹⁰⁸ accidental spills,¹⁰⁹ or ranges over a spatial gradient.¹¹⁰ High exposure concentrations may be used:¹¹¹ for assessing bioavailability in soil⁵⁸ in comparison to simpler media, or to accommodate analytical instrument detection limits. High concentrations are also used to establish no-effect limits, using limit tests; if an effect is not observed, the ENM is assumed to be non-toxic at lower concentrations,⁶² although effects could occur at longer exposure times. However, this approach is problematic if agglomeration and sedimentation of ENMs are concentration dependent, or if effect mechanisms do not scale with concentration, which might occur if organisms adaptively respond to toxicants.

Some challenges to ENM ecotoxicology are familiar to conventional chemical ecotoxicology while others are unique. With conventional chemicals, toxicity is related to the effective dose of a toxicant molecule crossing the cell membrane and disrupting essential processes. However, ENMs can exert effects as particulates (e.g., physical effects) and in a molecular or ionic form depending on the dissolution extent in the medium. The effective ENM dose—which impacts the assay results¹¹²—may be unknown or changing because it varies with media conditions or with dynamic physicochemical interactions of receptors and toxicants.¹¹³ One effective dose metric may be insufficient to characterize an

observed effect that could be related to both a physical interaction of the ENM particulate and the dissolved ionic form. Also, bioavailability and effective ENM doses¹¹⁴ change because ENMs can transform abiotically¹¹⁵ and biotically^{116, 117} during assessment. While such influences on stability exist in conventional ecotoxicology, stability in a particulate exposure must also consider the ENM size distribution, and potential physical changes to the ENM such as dissolution and agglomeration. Therefore, effective and nominal doses may or may not be related.

Although multiple ENMs may co-occur in product formulations, different ENMs are infrequently studied together¹¹⁸ or with co-contaminants.¹¹⁹⁻¹²¹ ENMs can conditionally sorb and modulate the toxicity and uptake of other contaminants and vice versa.¹²²⁻¹³⁴ ENMs can acquire coatings^{135, 136} such as natural organic matter (NOM),^{137, 138} and age with varying pH and sunlight.^{139, 140} Assessment outcomes are affected by media chemistry, physical characteristics, and additives (e.g., soil pH,¹⁴¹ temperature,¹⁴² or organic matter content,^{108, 143} and dispersing agents in aqueous media¹⁴⁴). The exposure concentration at the receptor and consequent effects¹⁴⁵ depend on ENM dissolution¹⁴⁶ perhaps assisted by biotic ligands,¹⁴⁷ and speciation,¹⁰³ shed ions,¹⁰³ surface associations,¹⁴⁸ and heteroaggregation with colloids and particulate matter,¹⁴⁸⁻¹⁵¹ or agglomeration and settling.¹⁵² All of these vary with ENM types and their varying properties, and ambient conditions.¹⁴⁸ Changes in ENM properties may change bioavailability^{94, 153} and toxicity.¹⁵⁴

Various test durations and endpoints have been studied, from acute responses measured by standard test protocols¹⁴¹ to short term microbial biodiversity and community composition effects¹⁰⁸ or plant genotoxicity^{111, 155} and nutrient composition changes.^{156, 157} Multiple endpoints, toxicant characterization methods, and experimental

controls increase assessment comprehensiveness¹⁵⁸ and reduce artifacts and misinterpretations.¹⁵² Experimental controls based on coatings and dispersants enable determining if the apparent toxicity is attributable to the ENM itself or to ligand or surface groups. Yet whether and how experimental controls are used varies widely, for example metal salts that allow interpreting biological responses relative to ENM dissolution products versus intact ENMs.^{159, 160}

The appropriate analytical method for quantifying, locating and characterizing organism-associated ENMs depends on ENM chemistry and amounts or concentrations. For ENMs exhibiting fluorescence or for ENMs containing heavy elements, high resolution microscopy can assess biological uptake and compartmentalization.^{99, 102, 161-163} X-ray synchrotron methods can sensitively locate bioaccumulated metal oxide ENMs¹⁶⁴ and their transformation products in biota.^{117, 165} No single analytical tool is suitable for all ENMs; however, the general lack of methodologies that can be routinely implemented to quantify ENM exposure in complex matrices continues to be a major challenge to the fundamental understanding of ecological effects.

Alternative testing strategies (e.g. miniaturized screening assessments using environmentally relevant bacteria)¹⁶⁶ can simultaneously assess many material types, controls, and concentrations,^{70, 75} which is useful for ENMs not available in sufficient quantities for microcosms. However, the potential for interferences in screening assay performance, depending on the specific ENM properties and toxicity test conditions,¹⁶⁷ are increasingly recognized, and therefore should be controlled or accounted for in study designs.^{152, 168}

To date, ENM ecological hazard assessments have not adequately explored numerous well-conceived and plausible exposure scenarios that are founded in theory, hypothesis, mechanism and occurrence probability; yet scenarios increase certainty and predictability when addressing nanotechnology-related material safety. For instance, hazard assessments have been mainly skewed towards as-produced ENMs without full consideration of potential aging, since aging cannot be fully standardized in a realistic context. However, such studies can and should further be conducted. Biological uptake, and the compartmentalization and speciation of ENMs, are still infrequently studied, limiting possibilities for attributing ENM exposure to effects at biological receptors.⁹ Varying degrees of rigor have been applied in designing and incorporating controls that are relevant to experimental questions or hypotheses. Varying degrees of attention are paid to experimental artifacts. While great advancements have been made in ENM ecotoxicology, improvements are needed to increase the environmental relevance of future research.

Recommendations

To understand hazards for plausible exposure initiation scenarios, assessment conditions will need to depart from standardized testing protocols.^{62, 169, 170} It may be helpful to link or compare data obtained for ENMs under plausible scenarios to those obtained with standard methods using standardized media (e.g., standard soil mixtures) to facilitate interpreting complex multivariate experiments or comparing results among multiple laboratories. Herein, several recommendations regard exposure conditions that should be used in assessing ENM ecotoxicity (Table 1).

Adequately characterize exposure conditions.

Soil ecotoxicity studies should specify the soil taxonomy¹⁷¹ and characteristics such as pH, clay content, and organic matter content, using standard methods.^{108, 172} Similarly, characterization of sediments should be provided. Natural soils and sediments are preferred, because artificial media do not harbor natural soil communities, and do not reflect chemical and physical characteristics (e.g. of specific surface area related to water holding, or of organic matter content) that influence ENM effects and bioavailability. Water characteristics determine ENM agglomeration, dissolution, and other behaviors affecting aquatic system compartmentalization (e.g., residing in the water column or depositing into sediments)⁹¹ and hence need definition. When adequately documented, media characteristics can be used retrospectively to interpret conditional ENM bioavailability or effect mechanisms.

Determine and report the form and concentrations of ENMs.

ENMs in testing should represent the particulate material form relevant to the environmental scenario (i.e., as-produced materials near manufacturing sites, or as-released with associated product surface coatings or with relevant co-contaminants in waste streams, including industrial byproducts, Fig. 1). ENMs should be fully characterized and their history (including storage conditions and time, or other uses that may affect compositional or physical characteristics) should be adequately described to allow comparing between studies. For toxic coatings,¹³⁶ coating identity and degree of coverage should be related to observed effects. Impurities in ENMs introduced during product synthesis and handling should be characterized, since they can sometimes account for

apparent ENM ecotoxicity.¹⁵² Nominal concentrations should scale according to exposure scenarios, or to specific objectives such as mechanistic research or quantifying biotic uptake. Dose verification, including size distribution and ENM concentration, is also desirable, although heretofore challenging in soil and sediment exposures.

ENM physicochemical changes during release and in the environment should be studied to uncover properties of ENMs that reach biological receptors. All potential forms of ENMs, including transformation products and residual reagents used in synthesis, should be accounted for, such that all toxicants can be related to biological responses.¹⁵² Study designs should anticipate dispersing agent effects¹⁴⁴ and the nature of transformed ENMs plus co-contaminants, by including controls to account for effects, fate, and kinetics of ENMs in the test medium.¹⁵⁴

The ENM physicochemical states should be understood before conducting hazard assessments.^{173, 174} This is important because some ENMs agglomerate or dissolve or otherwise change in laboratory media, resulting in non-uniform exposures and uncertain bioavailability. Such unevenness precludes relating measured effects to the applied dose. Spatial bio-association, bioaccumulation,¹⁷⁵ and intra-organism compartmentalization should be assessed (e.g., through imaging and mapping with sufficiently high resolution¹¹⁷) to locate ENMs and their components.

Relate biological receptors to the exposure scenario.

Important advances have been made toward characterizing the physicochemical factors influencing ENM behavior in environmental and test media, and towards utilizing that information to develop standardized methods for conducting ENM ecotoxicity

testing.^{62, 64, 67, 152} However, aquatic or terrestrial species^{8, 159, 166} —and even different species belonging to the same order¹⁷⁶ —respond differently to ENMs using the same tests. Thus, test species and endpoints should be carefully chosen to enhance the relevance of ENM ecotoxicity testing. Some complex interrelationships and dependencies between species comprising ecosystems have been described.^{8, 159, 177} However, focused research could rationally identify species for routine evaluations; likewise, the scientific rationale behind test species should be reported. Ecosystems are more complex than conditions of routine ENM ecotoxicity evaluations. Thus, research should define an optimal suite of test species and endpoints to determine the ecosystem response to a given ENM.

In general, biological receptors should be chosen for expected exposures stemming from realistic exposure scenarios. For example, relatively insoluble ENMs may, depending on their density, size and agglomeration state, rapidly settle out of suspension and associate with aquatic sediments. In that case, initial hazard assessment could focus on benthic, rather than pelagic, receptor organisms.⁶² Conversely, for ENMs that rapidly dissolve under environmental exposure conditions, conventional ecotoxicological exposure scenarios may be applied and receptors chosen to assess dissolution product toxicity. However, ENM dissolution rates vary, and pelagic organisms can be more sensitive than benthic organisms.^{100, 101} Thus, both ENM compartmentalization and form must be accounted for when choosing receptors.

Select endpoints relevant to appropriate receptor and exposure scenarios.

Multiple effects measurements¹⁵² should be applied to answer research questions. Rapid screening assessments should be prioritized within a testing strategy (Fig. 2).

Mechanistically understanding overt toxicity is needed, which may require measuring more omics⁸¹ endpoints (i.e., gene transcripts, proteins, and pathway metabolites) and choosing variables for developing mathematical models to predict toxicity at untested concentrations or conditions.⁸ Omics technologies can also identify potential modes of action (MOAs) that are conserved among different species. However, different scientific communities will have varying preferences in defining needs for omics-level investigations.

Perform adequate dosimetry to understand exposures.

Effects interpretation requires understanding the effective toxicant dose or other (perhaps indirect) basis of impacts.⁹ For ENMs, the mass concentration basis of dosing may relate only partially to the effective applied dose, since biological effects often originate from surface interactions with receptors.¹⁷⁸ Furthermore, ENMs are more complex than conventional chemicals because ENM shape, aggregation state and surface area may influence toxicity.^{67, 178-182} Thus, surface area applied has been suggested as a supplemental dosing metric.¹⁷⁸ However, ENM surface area in suspension/solid media is not a straightforward assessment given that ENMs may aggregate with a size distribution that is affected by the medium in which they are dispersed. In addition, coatings, either on pristine ENMs or acquired in the test media or environment, may alter toxicity.^{104, 183}

ENM amounts and forms effecting biological impacts should be understood and related to the administered dose to inform environmental risk assessment.^{9, 67, 184} This is the essence of dosimetry in ENM ecotoxicology. As with other exposure concerns related to hazard assessment, appropriate dose measurement depends on receptor and ENM characteristics, which are scenario-dependent. For example, mammalian cells are harmed

by ENMs that become internalized, yet uptake pathways depend on ENM characteristics.¹⁸⁵ Then again, bacterial receptors that affect ecosystem-level processes may be impacted by externally-associated ENMs at the cell membrane, or even in the surrounding environment. In those cases, dosimetry relies on understanding ENM behavior in the complex media in which bacteria reside (e.g., soil, water, sediments, and extracellular polymers^{186, 187}), which is scenario-driven. Endpoint observations of ENM damage will also depend on ENM processing in cells. During hazard assessments, understanding the history of biological interactions with internalized, or otherwise associated ENMs may not be feasible. Yet efforts should be made to measure and spatially associate ENM bioburden (quantity and form) within biological receptors, and to examine the relationships of applied ENMs to apparent effective dose and to effects.

Summary

Overall, it is not recommended to categorically exclude select conditions, environmental compartments, protocols, receptors, or endpoints, since any may be environmentally relevant. Rather, careful experimental designs around well-conceived, plausible exposure scenarios should be emphasized; also, ENM characteristics that influence biological responses under the dynamic conditions that occur in the environment and in biota should be characterized and quantified. One could imagine identifying key material-environment system determinants that could be systematically varied to provide test results across relevant determinant ranges. Such ideas are not specific to ENM ecotoxicology, but could establish defensible practices for making progress in hazard assessment while the ENM industry rapidly advances.

What exposure and design considerations drive mesocosm assessment of ENM environmental hazards?

Findings

Mesocosms are “enclosed experimental systems [that] are intended to serve as miniaturized worlds for studying ecological processes.”⁸⁴ While the distinctions between mesocosms and other experimental systems are not well delineated, mesocosms are generally larger experimental units and inherently more complex than bench-top microcosms or more simplified laboratory experiments.^{84, 188-191} Mesocosms for ENM ecotoxicology are intended to increase the complexity of experimental systems, such that more realistic ENM physical compartmentalization, speciation,¹¹⁷ and uptake into biota^{58, 192} can be achieved alongside biotic effects.^{89, 177} Also, the intent is to realistically characterize ENM fates and interactions with environmental system components, and to reveal fluxes among compartments of the ecosystems (e.g., the aqueous phase, sediments, and biota in aquatic systems) responsive to internal system influences that are unconstrained by investigator interventions.^{25, 177, 193}

Mesocosms have been used for testing relative biotic effects of ENM variants (e.g., surface functionalization, capping agents, adsorbed surface species or coatings),¹⁷⁷ and discerning ENM effects separately from effects of dissolution products (ions or complexes).¹⁹⁴ Mesocosm testing may occur following individual organism and microcosm studies (Fig. 2). For example, to study how ENMs impact crops, one could first establish the potential for hydroponic plant population impacts,¹¹¹ use soil microcosms to understand ENM bioavailability via observing soil microbial community shifts,^{85, 93} and then scale up to

greenhouse mesocosms of soil-grown crops. This sequence could provide an understanding of plant-microbe interactions,^{58, 87} ENM transformation and uptake in plants,¹¹⁷ and effects on food nutritional quality.¹⁵⁷ Still, there are relatively few published studies using mesocosms to assess ENM ecological hazards,⁵³ and the design and operating variables (including size, media, biota, length of study, and ENMs tested) of existing mesocosm studies are wide ranging.⁶⁰

By contrast, wastewater-associated ENMs,^{195, 196} and their transformations,¹⁹⁷⁻¹⁹⁹ effects, and fates²⁰⁰ in wastewater treatment plants (WWTPs),^{201, 202} along with the potential for ENMs to impact WWTP processes,^{48, 203, 204} have been more extensively studied. Since sewage contains ENMs, WWTPs are inherent forms of mesocosms.^{195, 205, 206} Studies at entire WWTP scales elucidate ENM fates during wastewater treatment, including significant association with biological treatment biomass^{201, 207} that becomes biosolids.^{76, 208} However, only 50% of biosolids produced in the U.S. are land-applied, and these biosolids are used on less than 1% of agricultural land in the U.S. (<http://www.epa.gov/biosolids>). Biosolids are land-applied even less in the European Union.²⁰⁹ Thus, knowledge of ENM fates in WWTPs and how final residues are disposed regionally are needed to develop plausible exposure scenarios.

Concerns with mesocosms include factors that can be difficult to control (e.g., pH or oxygenation) and that mesocosms may respond to artifacts including “wall” or “bottle” effects.⁸⁴ Further, mesocosms can conflate direct and indirect toxicant effects, typically do not have a full complement of control conditions, and deliver inconclusive results (e.g., where bioaccumulation may be explicable by either direct toxicant uptake or by trophic transfer). Biological communities in mesocosms also lack realistic ecological

interconnections, interactions, and energy flows. Nevertheless, outcomes can be improved by using carefully designed mesocosms and associated experiments.⁸⁴ For example, combined with analyzing mesocosm samples, performing practical “functional assays”³⁸ such as for heteroaggregation,²¹⁰ allows for anticipating phenomena and later interpreting ENM transformation and compartmentalization in mesocosms.¹⁷⁷ Similarly, batch physical association experiments—if conducted using realistic components, and over time frames that allow for quantifiable mass transfer—can assess ENM biomass association and readily suggest ENM fates in WWTPs.²⁰⁷ Still, hydrodynamic conditions are different in simplified tests versus mesocosms, which are different from those in the natural environment. Hydrodynamic conditions will impact ENM fate and transport and thus exposure concentrations at receptor boundaries. The inability to capture real environmental hydrodynamic conditions in any experimental scale is a general shortcoming for both ecotoxicology and transport studies.

Recommendations

Although mesocosms do not fully simulate real environments,⁸⁴ mesocosms are useful and should be employed, albeit judiciously due to their resource intensity, within a strategy (Fig. 2). Recommendations regarding using mesocosms for assessing ENM environmental hazards are provided in Table 2.

Mesocosm studies must be designed and conducted around well-conceived questions related to plausible exposure scenarios; they should use select endpoints, potentially including sensitive omics measurements,²¹¹ to answer questions or test hypotheses.⁸⁴ Internal process and constituent characterization should be thorough and

equally responsive to well-conceived, realistic scenarios. Functional assays, i.e. “intermediary, semi-empirical measures of processes or functions within a specified system that bridge the gap between nanomaterial properties and potential outcomes in complex systems”,³⁸ should precede mesocosm designs and experiments, and aid interpreting mesocosm results (Fig. 2).

Mesocosm artifacts are avoidable by following best practices for design and operation, although possible interferences of particulate material testing with assays must be evaluated.⁸⁴ As for other hazard assessments, ENMs should be tested across the product life cycle, (i.e., ranging from as-produced to released from products) within a motivating exposure scenario. Similarly, suitable material controls should be used to test hypotheses regarding ENM-specific effects (Table 1).

The recommendations made regarding exposure conditions in assessing ENM hazard potentials for model organisms (including in microcosms) should be followed for mesocosm studies (Table 1). Additionally, mesocosm designs should incorporate exposure durations, which should be sufficiently long to address population growth, reproduction, bioaccumulation, trophic transfer, and possibly transgenerational effects. Sufficient measurements of ENM concentrations and time-dependent properties must be made for clear interpretations.

Key to successfully interpreting mesocosm studies is using validated methods for measuring ENMs in complex media. Measurements should include the size distribution, concentration and chemical composition of ENMs in the test system, including biological tissues,^{161, 212-214} over time.²¹⁵ In some cases, transformation products are inventoried thoroughly during long term field-relevant exposures.¹⁹³ Detection schemes require sample

preparation to assess *in situ* exposures (e.g., aqueous or solvent extractions) before quantitative analyses, or drying and embedding before visual confirmation by electron microscopy. The potential for artifact introduction should be recognized. Recovery methods continually develop, such as cloud point extraction for concentrating ENMs from aqueous matrices.^{216, 217}

Depending on the exposure scenario, *in situ* aging may be a study objective. However, it is important to define what ‘aging’ really means and the specific application domain, since ‘aging’ is a wide-ranging term and can be used in different contexts, making comparisons impossible. At least, studies should be undertaken over sufficiently long time frames (e.g., acknowledging time scales of soil processes^{141, 218}), which may include repeated ENM applications,¹⁰⁸ such that appropriate aging, i.e. time-dependent transformation under realistic conditions, could occur. Alternatively, pre-aged ENMs could be used. However pre-aging protocols (whether in the context of product use conditions or under environmental conditions) are not yet standardized and, while some convention could allow for comparing across studies, the appropriate aging protocol would depend on the envisioned exposure scenario.

Co-contaminants should be considered and potentially introduced into mesocosms, since some ENMs sorb, concentrate, and increase exposure to other contaminants.¹³⁰ Select endpoints should account for ENMs as chemosensitizers.^{119, 120, 219, 220} Also, mesocosm study designs should anticipate and plan for measuring secondary effects (e.g., rhizosphere microbial shifts that affect plants⁸⁷ and soil nutrient turnover).

In summary, while few mesocosms have been used in assessing ENM ecotoxicity and are also rare for conventional chemical testing, such systems potentially offer greater

realism. Still, mesocosm exposure and design considerations should derive from immediate environmental applicability. The value of mesocosms to ENM ecotoxicology can increase by following recommendations including: addressing context-dependent questions while using relevant endpoints; considering and minimizing artifacts; using realistic exposure durations; quantifying ENMs and their products; and considering ENM aging, co-contaminants, and secondary biological effects (Table 2). Further, it should be acknowledged that mesocosms do not fully recreate natural environmental complexity. For example, aquatic mesocosms do not recreate actual environmental hydrodynamic, or temperature cycling, conditions. Hydrodynamics can significantly impact ENM aggregation or heteroaggregation, and fate and transport (e.g., sedimentation, dissolution, etc.). Therefore, potential impacts on the resulting concentrations at the receptor boundaries should be considered.

What is the state of knowledge regarding ENM environmental exposure conditions, via measurements or modeling simulations?

Findings

ENM environmental exposure conditions herein refer to where, how much, and in what forms ENMs may occur in the environment. These are central issues for ecotoxicology of ENMs because they suggest test exposure scenarios^{14, 15, 18} in which ENMs could impact biological receptors within environmental compartments (e.g., soil, sediments, air, and water) influenced by various factors (e.g., pH, redox potential, aeration, temperature, and ionic strength). These issues also influence outcomes of key regulatory concern: persistence, bioaccumulation, and toxicity.

In the parlance of exposure science,^{221, 222} three knowledge domains inform ecotoxicology, including the ecotoxicology of ENMs⁶:

- 1) “far-field”²²³ “environmental science”²²² regarding contaminant release, transport, fate, and transformation external to biological receptors;
- 2) “near-field”²²³ exposures at the receptor that lead to biological outcomes, as affected by contaminant bioavailability, affinity, uptake, accumulation, and biological transformation; and
- 3) “at point” responses and outcomes that occur at the biological receptor and include toxicity (Fig. 3).

Discharges (locations, types, and amounts) underpin exposure scenarios,^{14, 15, 37, 224} are initiated by situational contaminant releases (Fig. 1), and are referred to as source terms. Mass balance-based multimedia simulations²²⁵ mathematically account for released contaminants as they are transported and exchanged between environmental media, where contaminants may be transformed and may ultimately concentrate, potentially with altered compositions and structures (Fig. 3).

Far-field exposure modeling approaches vary by question, the modeling purpose (e.g., regulatory compliance, priority setting, research, material design, or industrial use), the required spatial resolution (e.g., a site or region, and the need to resolve environmental heterogeneity), the temporal conditions (e.g., steady state, dynamic, or episodic), and the predictive accuracy required.^{56, 226} Material Flow Analysis (MFA), which is a type of life-cycle inventory analysis, has been advanced to track ENM flows through various use patterns into volumes released into broad environmental compartments,²²⁷ scaled to regional ENM concentrations that release via WWTPs to water, air, landfills, and soil.²²⁷

Such models estimate exposure concentrations in part via engineering assumptions and in part via heuristics (i.e. expert judgment regarding the potential amount released to various media).²²⁸ Also, such material flow analysis models depend on the underlying data (e.g. production and release estimates) which are not readily available, making it difficult to validate model results and potentially leading to inaccurate estimates.^{32, 228}

Multimedia models for ENMs^{32, 37, 228-230} can predict environmental concentrations based on sources of continuous, time-dependent, or episodic releases^{32, 228} and are similar to multimedia models that predict environmental concentrations of organic chemicals²³¹⁻²³³ and particle-associated organic chemicals.^{233, 234} For ENMs, predicting particle size distribution—as affected by particle dissolution, agglomeration, and settling—is desired for various spatial and temporal endpoints. For one integrated MFA and multimedia model (MendNano and LearNano), user-defined inputs are flexible around product use and ENM release throughout material life cycles.²²⁸ It is noted that although validation of multimedia models is a formidable task, various components of such models have been validated as well as model predictions with such models for particle-bound pollutants.

Most far-field²²³ models of ENMs have major challenges. First, the quantities and types of ENMs being manufactured are unknown to the general public due to issues surrounding confidential business information, leading to a reliance on market research.⁵³ The resulting public uncertainty will persist while nanotechnology continues a course of rapid innovation, as is typical of new industries.²³⁵ The rates of product use and ENM releases at all life-cycle stages (i.e., manufacturing, waste handling, product use, and product disposal) are also not defined.²²⁵ There are challenges associated with modeling transport processes through specific media and across media (i.e., water, soil, air), highly

divergent time scales of processes, lack of required input parameters, and the need for validation of results (e.g., through measurements of ENMs in the environment).²²⁶ Several multimedia models developed for conventional chemicals could be adapted around ENMs, but few account for fate processes specific to nanoparticles (e.g., never in thermodynamic equilibrium).^{37, 236} In addition, various transport models for a single medium and in the multimedia environment could be adapted for far-field analysis of ENMs, but few account for fate processes distinctive to ENMs (e.g., considering the complexity of ENMs aggregating and that their transport is governed by physical rather than thermodynamic driving forces).^{37, 236} Moreover, their validation, which would require ENM monitoring data, is a major challenge.

The lack of understanding of many fundamental ENM behaviors under environmental conditions propagates into broad uncertainties, for example in predicting ENM removal to solids or aqueous fractions in WWTPs.^{225, 227} ENM surface chemistries fundamentally affect ENM agglomeration or dispersion²³⁷ and likely affect bioavailability.⁹⁴ Some species on ENM surfaces may degrade in the environment,²³⁸ while other adsorbates can be acquired.^{239, 240} Carbonaceous ENMs may be transformed or degraded by environmental processes such as photo-,^{241, 242} enzymatic,^{243, 244} chemical,²⁴⁵ and bio-degradation.²⁴⁶ Redox and other environmental conditions will affect nanomaterial surfaces, which for nano-Ag includes formation of sulfide that inhibits dissolution.^{39, 247} Surface chemistry also affects transformation rates of primary particles and aggregates (e.g., in WWTP biological processes).²⁴⁸ For many ENMs such as nano-ceria,²⁴⁹ reactivity is highly size-dependent. To accurately model material fates thus requires understanding how material surface properties affect integrity, how both change under varying

environmental conditions such as pH, clay content,²⁵⁰ and organic matter content,^{143, 251} and how surface properties and particle reactivity affect physicochemical processes that are parameterized in far-field models. This is especially true for ENMs used as pesticide delivery mechanisms, including carbon nanotube composites with specifically-reactive surface monomers. Yet only recently has modeling attempted to address differing properties of a material's structural variants (e.g., photocatalytic versus photostable nano-TiO₂).²³⁰

Evaluating computational model predictions is a challenge for ENMs, which presently are estimated to occur in the environment at low (ng/L, to mg/L) concentrations.^{32, 230, 252-254} Also, detection methods for ENMs in environmental media and distinguishing ENMs from natural chemical analogs are still under development,^{161, 255-257} with more evaluation strategies needed including a framework for validating new ENM analytical detection methods.²¹⁵ Fullerenes from incidental sources (combustion products) were quantified in river sediments collected from locations across the globe²⁵⁸ and quantified in the atmosphere over the Mediterranean Sea.²⁵⁹ Perhaps related to a viable exposure scenario, fullerenes were quantified at relatively high concentrations in treated wastewater effluent²⁶⁰ and at ng/L to µg/L²⁶¹ concentrations in river waters receiving effluent discharge. While not necessarily nanoscale, similarly high concentrations of TiO₂ were reported for sediments sampled near a WWTP outfall.²⁶²

The greatest uncertainty in ENM exposures is near-field (Fig. 3), at the receptor where toxicant dose manifests as internal dose. Heteroaggregation is a dominant fate process for ENMs when they interact with natural colloids.^{37, 263-265} Given sufficient residence time for ENMs in environmental matrices, heteroaggregation (i.e., kinetically-

controlled association with colloids) and to a lesser degree homoaggregation will affect localized compartmentalization, including stability in the water column and therefore, sedimentation.¹⁷⁷ However, these processes do not preclude biological impacts under simulated environmental conditions, as has been shown for nano-ceria in a complex aquatic mesocosm.¹⁷⁷ Exposure can be confirmed by quantifying receptor body burdens, thereby allowing for quantitatively relating near-field exposure to biological effects.¹⁷⁷ Thus, in the absence of detailed, biologically complex, near-field models for local exposures to environmental receptors, the ability to trace ENMs to biological receptors sampled directly from the environment becomes the best available approach to relate far-field exposures to biological impacts.¹⁷⁷

Overall, material flow models and multimedia modeling of ENMs have advanced to inform ENM ecotoxicology. Available far-field modeling frameworks are adaptable to changing inputs despite uncertainties in production volumes. Major uncertainties remain at the nexus of ENM surface and core chemistries as related to nanomaterial transport, aggregation, and degradation characteristics. However, fundamental research (e.g., of conditional hetero- and homoaggregation, and of dissolution) is needed to discover and parameterize complex fate processes. New approaches, such as assays that can be used to rapidly probe surface associations,³⁸ demonstrate how to populate far-field models and how to determine near-field exposures associated with effects. Although existing models can simulate particle movement, deposition, and some transformations, the knowledge state regarding ENM environmental exposure conditions via measurements or modeling simulations cannot be assumed to accurately represent actual conditions at biological receptors.

680

681 Recommendations

682 Despite significant progress in modeling potential ENM environmental distributions,
683 several key constraints exist. Models predict the average concentration within broad
684 compartments such as soils, sediments, water, or wastewater, even when constrained to
685 regional releases. Actual concentrations at biological exposure sites are typically unknown,
686 but can be constrained within upper and lower bounds. Due to the potential for bio-
687 association, bioaccumulation, and hetero- or homoaggregation, predicted environmental
688 ENM concentrations may inaccurately estimate bioactive concentrations at receptors.
689 Following are recommendations for using measurements or modeling simulations to
690 understand conditions of ENM environmental exposures.

691 • The limitations of modeling approaches should be recognized more broadly, such that
692 generic model predictions are not used to rigidly define or judge ecotoxicity testing
693 conditions.

694 • Modeling should be simplified where possible and coupled with evaluation and
695 iteration while varying assumptions to develop an understanding of the sensitivity of
696 modeling parameters.

697 • Meteorological conditions play an important role in ENM transport, and thus should be
698 carefully documented when assessing model predictions relative to monitoring.

699 • ENM surface characteristics relevant to the receptor and the exposure context should
700 be considered by modelers and experimentalists. Considerations should include: which
701 ENM characteristics prevail near receptors, how this changes in the environment, and
702 how particle stability, migration and bioavailability are affected.

- Batch experimental approaches to rapidly assess relationships between environmental factors (e.g., pH, redox potential or aeration, ionic strength, and organic matter content) and ENM physicochemical characteristics that affect fate processes (e.g., homo- and heteroaggregation, sedimentation, and dissolution) could be more routinely incorporated into assessments (Fig. 2), to justify further testing and for interpreting other testing results.
- Improvements in quantifying ENMs in complex media, including biological tissue, are needed to validate multimedia models and to determine near-field exposures to biological receptors.
- Improved fundamental understanding of how ENM variants (e.g., in transformable coatings, shapes, sizes, or primary chemistry) comparatively migrate under varying environmental conditions can be advanced through research that intentionally develops and uses relatively stable ENMs as tracers.^{188, 189, 266}

In summary, understanding exposure conditions should continue to improve from fundamental experimental research of ENM environmental behaviors. Modeling should continue to improve with such understanding, and avenues for model evaluation should continue to grow. Environmental relevance in ENM ecotoxicology need not be defined or constrained by model outputs, especially given recognized uncertainties and the need for continued model evolution. Lastly, echoing prior recommendations,¹⁵ the efforts of exposure modelers, analytical methods developers, and ecotoxicologists should continue to influence each other for maximum individual and mutual benefit.

How should concepts such as exposure conditions, ENM transformation, dose, and body burden be utilized in interpreting biological and computational findings for assessing risks?

Findings

Many of the outstanding research issues and recommendations for evolving ENM ecotoxicology (Tables 1 and 2) are echoed in the discourse for other chemicals of emerging concern (CECs).²⁶⁷ These include the need for systematically understanding ENM and decomposition product toxicity across various receptors within linked levels of biological organization,^{8, 268, 269} quantifying actual exposures²⁷⁰ and uptake²⁷¹ into environmental receptors,^{57, 161} gaining mechanistic insights into and biological markers for acute and chronic low level exposures,^{272, 273} and understanding how environmental factors including co-contaminants affect ENM transformation²⁷⁴ and biological impacts. Still, how can the potential for exposure and impacts of ENMs be anticipated, prevented, managed, or mitigated? Further, what data and tools do decision makers need to inform their work? Innovation in nanotechnology hinges on having the science to evaluate ENM safety.

While no formalized process for incorporating all exposure conditions and concepts of ENM transformation, dose, and body burden into risk assessments currently exists, a proposed framework approach to risk characterization over the life cycle of ENMs has been published and is available.⁶³ This framework advocates an initial decision cut-off in regards to exposure; in the absence of exposure, the need for further assessment is diminished or negated.^{26, 63} In this available framework, ENMs that are certain to rapidly dissolve into ionic components in a destined environmental compartment would be assessed for risk based on the released components rather than the original nanoparticles.⁶³ Persistent

ENMs are expected to accumulate in matrices such as sediments.⁶³ The consequences of ENMs to successive generations, biodiversity, and ecosystem services⁴⁹ are not addressed by model organism-specific assays of discrete growth and mortality.^{8, 63} Nonetheless, in this available framework, toxicity endpoints associated with standardized testing protocols for sediment, aquatic, and terrestrial standard population-level endpoints over short and long time frames are advocated for assessing hazards of simulated ENM concentrations in the environment.⁶³ In this framework, sunlight is an environmental variable, bioaccumulation is measured, and ENM modifications during product and material life cycles that may change bioavailability are considered.⁶³ While such a framework has broad organizational appeal, priority setting within the framework is required and thus could focus on tests that are relatively well aligned with likely exposure scenarios.

Even with a risk assessment framework that considers ENMs across product life cycles and considers sediments, water, and soil in testing endpoints,⁶³ major hurdles hinder regulatory agencies, and research scientists, in using concepts such as exposure conditions, ENM transformation, dose, and body burden in interpreting biological and computational findings for assessing risks. Toxicity tests developed for dissolved chemicals typically require significant modification for use with ENMs.⁵⁷ Tests may not apply to ENMs if they are not appropriate for solids.²⁷⁵ Additional scientifically-based hazard information from the peer-reviewed literature may or may not be available for consideration. ENMs used in ecotoxicity tests, which are sometimes laboratory-synthesized to overcome uncertainty regarding proprietary coating or other commercial formulations, may be insufficiently analogous to allow for extrapolating information or risk comparisons.

Issues include the need to know test material characteristics and how they relate to testing results and the ENM life cycle. Even if an initial risk assessment considers ENM solubility,⁶³ ENM dissolution is not instantaneous; therefore, at what stage of dissolution does the contaminant no longer pose a hazard as an ENM? Also, where biological impacts stem from ENM surface characteristics, how can mass concentration be used to judge hazards? Environmental ENM effects in benchtop experiments can be indirect, stemming from physical nutrient depletion,¹⁵² or amplifying organism uptake of co-contaminants.²⁷⁶ Other indirect physical effects derive from ENMs adhering to the organism surface,²⁷⁷ light shading,²⁷⁸ or internal food displacement.⁹⁹ Near-field exposures (Fig. 3) can result in biological hazards from specific ENMs based on their properties (e.g., electronic).^{75, 279}

By definition, ecological risk assessment (ERA) is “the process for evaluating how likely the environment will be impacted as a result of exposure to one or more environmental stressors.”²⁸⁰ ERA involves predicting effects for individuals, populations, communities and ecosystems, and concerns itself with valuable⁴⁹ ecosystem services such as nutrient cycling.²⁸⁰ Thus, conducting ERAs for ENMs could benefit from an ecological outlook. All levels of biological organization, and interactions between them, would be considered when assessing responses to ENM exposure (Fig. 3).

Release and exposure scenarios (Fig. 1), use of functional assays for assessing environmental compartmentalization (Fig. 2),³⁸ and combined life-cycle and multi-media modeling²²⁸ have important roles in focusing ENM ecotoxicology. Less recognized is that mechanistically based models of dynamic biological effects are informed by hazard assessment research. Different types of process-based, dynamic models (e.g., Adverse Outcome Pathway or AOP,^{281, 282} and Dynamic Energy Budget or DEB²⁸³) allow for

793 predicting effects from exposures stepwise, starting at subcellular levels, into individuals,
794 through populations, and conceivably to communities and ecosystems. Developing process-
795 based models requires researching key effects processes^{71, 284, 285} and ecological
796 feedbacks.¹⁵³ Models are formalized to describe interactive processes culminating in
797 toxicity such as reactive oxygen species (ROS) generation and cellular damage. Process-
798 based mathematical expressions evolve with empirically-based discoveries or through
799 model reconciliation with experimental data. Parameters are independent of toxicity
800 testing protocols, although models could be informed by standard test results. Thus, ENM
801 ecotoxicity research could support predictive toxicology by informing and populating
802 process-based, dynamic ecological effects models.

803 A comprehensive fate and effects research agenda is needed for addressing ENM
804 quantification in complex media.^{286, 287} Such an agenda has allowed for assessing
805 experimental compartmentalization,²⁸⁸ and sensitively assessing environmental
806 persistence,²⁸⁹ toxicity, bioaccumulation,^{287, 290} trophic transfer,²⁹¹ and indirect effects from
807 the uptake of ENMs coated in other hazardous materials.^{292, 293} Such research could
808 substantially inform ENM risk assessment for a relevant environmental exposure scenario.
809 However, most ENMs have not been studied comprehensively along the entire exposure
810 and effects continuum (Fig. 3). Further, the approach is not sustainable. Rather, the need is
811 to develop efficient approaches applicable within an overall approach to rapidly evaluate
812 the large number of ENMs under commercialization (Fig. 2). A research agenda that focuses
813 on distilling key determinants of exposure and hazard for ENM-environment systems that
814 can be measured experimentally would be most compelling.

Thus, while the science of ENM ecotoxicology and exposure characterization has advanced, there are disconnects between how regulators review ENM-based products for environmental safety and the research that is conducted to evaluate hazards. Except for results published in open source outlets or directly reported, research may be unknown to government bodies. Ongoing synthesis of published research results is challenging due to high variability across study conditions and ENMs tested, and due to effort needed to regularly update such comparisons. Moreover, there is a systematic resistance to publishing “no effect” studies in the peer-reviewed literature.²⁹⁴ As a result, relying only on published research to inform regulatory decisions can present challenges. A life cycle-based framework facilitates exposure modeling and hazard testing to support risk assessment. However, extrapolation of effects to untested concentrations, study, or environmental conditions, and across biological levels of organization, requires understanding dynamic biological process-based effects, which current standard tests neither deliver nor sufficiently inform. Ultimately, exposure scenarios are useful for framing and focusing ENM ecotoxicology, and some version of a tiered intelligent testing and risk assessment (Fig. 2) strategy is needed.

Such a conceptual tiered strategy considering the impact of the ENMs’ varying properties on ecological risks at different life cycle stages was proposed in the EU FP7 MARINA (Managing Risks of Nanoparticles) project²⁹⁵ and is being further developed in the EU NANoREG programme. This strategy considers several domains (exposure, fate, kinetics, and hazard) represented by specific tools ranging from relatively simple in the lower tiers to more complex and specific in the higher tiers. The framework aim is to structure information collection and generation for cost-efficient risk assessment,

compliant with 3R animal-use testing principles (i.e., replacement, reduction, and refinement), which should also be pursued by means of grouping ENMs.

A strategy for grouping ENMs based on releases, uses, physicochemical properties, bioaccumulation, bioavailability, and effects for both human and ecological risk assessment is currently in development across a number of EU research projects such as MARINA, NANoREG, SUN, and GUIDEnano. These efforts have been challenged by the complexity of ENM identity and interactions, but this approach is necessary, as the costs for safety assessment on a case-by-case basis would be exorbitant.²⁹⁵ Therefore, a vision on ENM grouping is needed, which should apply in a regulatory context.^{296, 297} Applying grouping in regulatory risk assessments should enable read-across, i.e., filling a data gap by using information on one ENM, or a non-ENM, for another substance in the same group.

Recommendations

The abovementioned tools should be fitted into a risk assessment strategy for ENMs. This strategy should be flexible enough to address different assessment goals depending on the user's needs, considering all data already available as a starting point, contingent upon data quality evaluation²⁹⁸ and selecting the most appropriate tools to fill existing data gaps. Such a strategy should ideally be exposure-driven, starting with identifying the most relevant exposure scenarios in the ENM life cycle, and evaluating completeness and quality of the available data from a risk assessment perspective. This facilitates careful prioritization of ENMs to optimize testing efforts and can inform more realistic ecotoxicological investigations. Doing so can allow one to screen-out irrelevant exposure routes, eliminate unnecessary testing, and support prioritization of exposure scenarios.

Exposure assessment should begin with an analysis of plausible exposure scenarios; where none is expected, further testing may be precluded for the applicable use patterns and volumes.^{26, 63} Researchers and regulators need to understand actual exposures at biological receptors. This exposure-driven approach can also provide important information on realistic environmental conditions to affect test designs for improved interpretation of laboratory toxicology studies. Such practices can ensue in the interim, while research continues to discover best hazard assessment practices.

Experimental ENM toxicity assessments, using ecologically relevant receptors and across linked biological levels of organization, should inform developing and parameterizing dynamic process-based models. Such models should respond to future scenarios and predict impacts. ENM characteristics, exposure conditions, and ENM transformation, dose, and body burden should be used in interpreting biological and computational findings for assessing ENM risks. ENM test results should be benchmarked to results for appropriate controls to establish relative hazard (e.g., comparing dissolving ENMs to their dissolution products or comparing persistent ENMs to non-nano analogs). This applies to pinnacle concerns in ecological fate assessment of bioaccumulation, biomagnification, and biopersistence.²⁹⁹

How to develop, interpret, and use pertinent information in ENM environmental risk assessment is a larger issue that should become part of an extended dialog among regulators, industry, civil society organizations, researchers, and other societal members so that the fundamental research will inform decision making. Collaborative decisions are recommended for focusing ENM ecotoxicology towards relevant scenarios, including

testing the most relevant materials throughout ENM life cycles and employing appropriate hazard assessment approaches, towards meaningful ecological risk assessment.

Summary and Outcomes: Towards Defining “Environmental Relevance”

The overarching question motivating this critical review was: how can we ensure that hazard assessment in ENM ecotoxicology is as environmentally relevant as possible? The answer requires considering how ecotoxicity tests are performed, what constitutes pertinent concentration and test conditions for ENMs (Table 1), the main biotic and abiotic attributes of the environment, how ecologically oriented hazard assessment is undertaken (Table 2), and how the resulting information should be interpreted. Answering this question yielded three primary insights. First, environmental relevance is informed by a logical consideration of what exposures might occur, to which receptors, and to what outcomes. The consideration should begin with a plausible release and exposure scenario (Fig. 1), and use best available knowledge and technologies to develop the full assessment approach. Concerns regarding ENM concentrations used in hazard assessments are paramount, but are not the only concerns. ENM concentrations should be selected to assess potential effects, but overly high concentrations that fundamentally change media conditions should be avoided. Still, concentrations ranging above and below predicted ENM average concentrations must be assessed for understanding potential organismal effects, underlying mechanisms and their concentration dependencies, and for informing process-based dynamic biological effects models. In addition to the nanomaterial, the conventional material (e.g., dissolved metal) should be tested. ENM distributions and fates in broad environmental compartments do not equate to concentrations and forms near, or effective

at, actual biological receptors.¹¹⁴ Therefore, research results on ENM effects should not be disregarded on the limited basis of environmentally relevant exposure concentrations when the study conditions were predicated on a broader hypothesis.

In addition to tethering ENM ecotoxicology to exposure initiation scenarios (Fig. 1), the concept of employing tiered approaches in hazard and risk assessment resonated (Fig. 2). Multi-stage approaches to ENM hazard assessment are advocated.¹¹⁴ A highly developed tiered approach for health and safety testing of nanotechnologies has been published²⁶ and strategies for tiered risk assessment and grouping are underway.^{295, 297, 298} Staging ENM ecotoxicology efforts, such that potential interactive impacts at all levels of biological organization (Fig. 2 and 3) are evaluated, could simultaneously inform risk assessment and predictive process-based effects model development. As some ENMs can cause biological impacts from ENM properties or characteristics,³⁰⁰ ENM ecotoxicology should be oriented to logical exposure initiation scenarios (Fig. 1) based on ENM life cycles, via testing tiers (Fig. 2). Finally, coordination is recommended among multiple disciplines in ENM environmental analysis, fate and transport modeling, and hazard assessment, towards rapidly advancing research using tiered approaches around realistic exposure scenarios.

Acknowledgements

Funding was from the National Science Foundation and the U.S. Environmental Protection Agency under Cooperative Agreement DBI-1266377. Any opinions, findings, and conclusions are those of the authors and do not necessarily reflect the views of the National Science Foundation, the U.S. Environmental Protection Agency, or the National Institute of Standards and Technology. Although this document has been reviewed in accordance with

929 U.S. Environmental Protection Agency policy and approved for publication, the views are
930 those of the authors and may not necessarily reflect official Agency policy. Mention of trade
931 names or commercial products should not be interpreted as endorsement or
932 recommendation for use. The authors acknowledge the helpful comments of three
933 anonymous reviewers.

934

935 **Supporting Information Available**

936 Supporting Information includes a description of the process and methods used to develop
937 this critical review, one supporting table, and one supporting figure.

References

1. Renn, O.; Roco, M. C., Nanotechnology and the need for risk governance. *Journal of Nanoparticle Research* **2006**, *8*, (2), 153-191.
2. Roco, M. C., The long view of nanotechnology development: the National Nanotechnology Initiative at 10 years. *J. Nanopart. Res.* **2011**, *13*, (2), 427-445.
3. Schaumann, G. E.; Baumann, T.; Lang, F.; Metreveli, G.; Vogel, H.-J., Engineered nanoparticles in soils and waters. *Science of The Total Environment* **2015**, *535*, 1-2.
4. U. S. Environmental Protection Agency, Nanotechnology White Paper. In Council, S. P., Ed. U.S. Environmental Protection Agency: 2007; p 120.
5. The Royal Society *Nanoscience and nanotechnologies: opportunities and uncertainties*; The Royal Society and The Royal Academy of Engineering: London, 2004; p 116.
6. Klaine, S. J.; Alvarez, P. J. J.; Batley, G. E.; Fernandes, T. F.; Handy, R. D.; Lyon, D. Y.; Mahendra, S.; McLaughlin, M. J.; Lead, J. R., Nanomaterials in the environment: behavior, fate, bioavailability, and effects. *Environmental Toxicology and Chemistry* **2008**, *27*, (9), 1825-1851.
7. Stark, W. J.; Stoessel, P. R.; Wohlleben, W.; Hafner, A., Industrial applications of nanoparticles. *Chemical Society Reviews* **2015**, *44*, (16), 5793-5805.
8. Holden, P. A.; Nisbet, R. M.; Lenihan, H. S.; Miller, R. J.; Cherr, G. N.; Schimel, J. P.; Gardea-Torresdey, J. L., Ecological nanotoxicology: integrating nanomaterial hazard considerations across the subcellular, population, community, and ecosystems levels. *Accounts of Chemical Research* **2013**, *46*, (3), 813-822.
9. Hinderliter, P.; Minard, K.; Orr, G.; Chrisler, W.; Thrall, B.; Pounds, J.; Teeguarden, J., ISDD: A computational model of particle sedimentation, diffusion and target cell dosimetry for in vitro toxicity studies. *Particle and Fibre Toxicology* **2010**, *7*, (1), 36.
10. Mudunkotuwa, I. A.; Pettibone, J. M.; Grassian, V. H., Environmental implications of nanoparticle aging in the processing and fate of copper-based nanomaterials. *Environmental Science & Technology* **2012**, *46*, (13), 7001-7010.
11. Borm, P.; Klaessig, F. C.; Landry, T. D.; Moudgil, B.; Pauluhn, J.; Thomas, K.; Trottier, R.; Wood, S., Research strategies for safety evaluation of nanomaterials, Part V: Role of dissolution in biological fate and effects of nanoscale particles. *Toxicological Sciences* **2006**, *90*, (1), 23-32.
12. Burello, E.; Worth, A. P., A theoretical framework for predicting the oxidative stress potential of oxide nanoparticles. *Nanotoxicology* **2010**, *0*, (0), 1-8.
13. Nel, A. E.; Madler, L.; Velegol, D.; Xia, T.; Hoek, E. M. V.; Somasundaran, P.; Klaessig, F.; Castranova, V.; Thompson, M., Understanding biophysicochemical interactions at the nano-bio interface. *Nature Materials* **2009**, *8*, (7), 543-557.
14. Nowack, B.; Ranville, J. F.; Diamond, S.; Gallego-Urrea, J. A.; Metcalfe, C.; Rose, J.; Horne, N.; Koelmans, A. A.; Klaine, S. J., Potential scenarios for nanomaterial release and subsequent alteration in the environment. *Environmental Toxicology and Chemistry* **2012**, *31*, (1), 50-59.

15. Praetorius, A.; Arvidsson, R.; Molander, S.; Scheringer, M., Facing complexity through informed simplifications: a research agenda for aquatic exposure assessment of nanoparticles. *Environmental Science: Processes & Impacts* **2013**, *15*, (1), 161-168.
16. Eckelman, M. J.; Mauter, M. S.; Isaacs, J. A.; Elimelech, M., New perspectives on nanomaterial aquatic ecotoxicity: production impacts exceed direct exposure impacts for carbon nanotubes. *Environmental Science & Technology* **2012**, *46*, (5), 2902-2910.
17. Walser, T.; Demou, E.; Lang, D. J.; Hellweg, S., Prospective environmental life cycle assessment of nanosilver t-shirts. *Environmental Science & Technology* **2011**, *45*, (10), 4570-4578.
18. Gottschalk, F.; Ort, C.; Scholz, R. W.; Nowack, B., Engineered nanomaterials in rivers - exposure scenarios for Switzerland at high spatial and temporal resolution. *Environmental Pollution* **2011**, *159*, (12), 3439-3445.
19. Aitken, R. J.; Chaudhry, M. Q.; Boxall, A. B. A.; Hull, M., Manufacture and use of nanomaterials: current status in the UK and global trends. *Occupational Medicine-Oxford* **2006**, *56*, (5), 300-306.
20. Hong, J.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Nanomaterials in agricultural production: benefits and possible threats? . In *Sustainable Nanotechnology and the Environment: Advances and Achievements*, Shamin, N.; Sharma, V. K., Eds. American Chemical Society: 2013; Vol. 1124, pp 73-90.
21. MacCuspie, R. I.; Hyman, H.; Yakymyshyn, C.; Srinivasan, S. S.; Dhau, J.; Drake, C., A framework for identifying performance targets for sustainable nanomaterials. *Sustainable Materials and Technologies* **2014**, *1-2*, 17-25.
22. Johnson, A. C.; Park, B., Predicting contamination by the fuel additive cerium oxide engineered nanoparticles within the United Kingdom and the associated risks. *Environmental Toxicology and Chemistry* **2012**, *31*, (11), 2582-2587.
23. Arvidsson, R.; Molander, S.; Sanden, B. A., Impacts of a silver-coated future. *Journal of Industrial Ecology* **2011**, *15*, (6), 844-854.
24. Wigger, H.; Hackmann, S.; Zimmermann, T.; Köser, J.; Thöming, J.; von Gleich, A., Influences of use activities and waste management on environmental releases of engineered nanomaterials. *Science of The Total Environment* **2015**, *535*, 160-171.
25. Colman, B. P.; Arnaout, C. L.; Anciaux, S.; Gunsch, C. K.; Hochella, M. F., Jr.; Kim, B.; Lowry, G. V.; McGill, B. M.; Reinsch, B. C.; Richardson, C. J.; Unrine, J. M.; Wright, J. P.; Yin, L.; Bernhardt, E. S., Low concentrations of silver nanoparticles in biosolids cause adverse ecosystem responses under realistic field scenario. *PLoS ONE* **2013**, *8*, (2), e57189.
26. Collier, Z. A.; Kennedy, A. J.; Poda, A. R.; Cuddy, M. F.; Moser, R. D.; MacCuspie, R. I.; Harmon, A.; Plourde, K.; Haines, C. D.; Steevens, J. A., Tiered guidance for risk-informed environmental health and safety testing of nanotechnologies. *Journal of Nanoparticle Research* **2015**, *17*, (3), 1-21.
27. Windler, L.; Lorenz, C.; von Goetz, N.; Hungerbühler, K.; Amberg, M.; Heuberger, M.; Nowack, B., Release of titanium dioxide from textiles during washing. *Environmental Science & Technology* **2012**, *46*, (15), 8181-8188.
28. Stacey, H.; Wendel, W.; Maria, D.; Bernd, N.; Shaun, C.; Richard, C.; Andrew, M., Measuring nanomaterial release from carbon nanotube composites: review of the state of the science. *Journal of Physics: Conference Series* **2015**, *617*, (1), 012026.

29. Nowack, B.; David, R. M.; Fissan, H.; Morris, H.; Shatkin, J. A.; Stintz, M.; Zepp, R.; Brouwer, D., Potential release scenarios for carbon nanotubes used in composites. *Environment International* **2013**, *59*, (0), 1-11.
30. Hirth, S.; Cena, L.; Cox, G.; Tomovic, Z.; Peters, T.; Wohlleben, W., Scenarios and methods that induce protruding or released CNTs after degradation of nanocomposite materials. *Journal of Nanoparticle Research* **2013**, *15*, (4), 1-15.
31. Petersen, E. J.; Lam, T.; Gorham, J. M.; Scott, K. C.; Long, C. J.; Stanley, D.; Sharma, R.; Liddle, J. A.; Pellegrin, B.; Nguyen, T., Methods to assess the impact of UV irradiation on the surface chemistry and structure of multiwall carbon nanotube epoxy nanocomposites. *Carbon* **2014**, *69*, 194-205.
32. Liu, H. H.; Cohen, Y., Multimedia environmental distribution of engineered nanomaterials. *Environmental Science & Technology* **2014**, *48*, (6), 3281-3292.
33. Goodwin, D. G.; Marsh, K. M.; Sosa, I. B.; Payne, J. B.; Gorham, J. M.; Bouwer, E. J.; Fairbrother, D. H., Interactions of microorganisms with polymer nanocomposite surfaces containing oxidized carbon nanotubes. *Environmental Science & Technology* **2015**, *49*, (9), 5484-5492.
34. Khan, F. R.; Laycock, A.; Dybowska, A.; Lerner, F.; Smith, B. D.; Rainbow, P. S.; Luoma, S. N.; Rehkaemper, M.; Valsami-Jones, E., Stable isotope tracer to determine uptake and efflux dynamics of ZnO nano- and bulk particles and dissolved Zn to an estuarine snail. *Environmental Science & Technology* **2013**, *47*, (15), 8532-8539.
35. Filser, J.; Arndt, D.; Baumann, J.; Geppert, M.; Hackmann, S.; Luther, E. M.; Pade, C.; Prenzel, K.; Wigger, H.; Arning, J.; Hohnholt, M. C.; Koser, J.; Kuck, A.; Lesnikov, E.; Neumann, J.; Schuttrumpf, S.; Warrelmann, J.; Baumer, M.; Dringen, R.; von Gleich, A.; Swiderek, P.; Thoming, J., Intrinsically green iron oxide nanoparticles? From synthesis via (eco-)toxicology to scenario modelling. *Nanoscale* **2013**, *5*, (3), 1034-1046.
36. Levard, C.; Hotze, E. M.; Lowry, G. V.; Brown, G. E., Environmental transformations of silver nanoparticles: impact on stability and toxicity. *Environmental Science & Technology* **2012**, *46*, (13), 6900-6914.
37. Praetorius, A.; Scheringer, M.; Hungerbuhler, K., Development of environmental fate models for engineered nanoparticles-a case study of TiO₂ nanoparticles in the Rhine River. *Environmental Science & Technology* **2012**, *46*, (12), 6705-6713.
38. Hendren, C. O.; Lowry, G. V.; Unrine, J. M.; Wiesner, M. R., A functional assay-based strategy for nanomaterial risk forecasting. *Science of the Total Environment* **2015**, *536*, 1029-1037.
39. Kaegi, R.; Voegelin, A.; Sinnet, B.; Zuleeg, S.; Siegrist, H.; Burkhardt, M., Transformation of AgCl nanoparticles in a sewer system — A field study. *Science of The Total Environment* **2015**, *535*, 20-27.
40. Liu, X. Y.; Vinson, D.; Abt, D.; Hurt, R. H.; Rand, D. M., Differential toxicity of carbon nanomaterials in *Drosophila*: larval dietary uptake is benign, but adult exposure causes locomotor impairment and mortality. *Environmental Science & Technology* **2009**, *43*, (16), 6357-6363.
41. Petersen, E. J.; Pinto, R. A.; Zhang, L. W.; Huang, Q. G.; Landrum, P. F.; Weber, W. J., Effects of polyethyleneimine-mediated functionalization of multi-walled carbon nanotubes on earthworm bioaccumulation and sorption by soils. *Environmental Science & Technology* **2011**, *45*, (8), 3718-3724.

42. Johnson, A. C.; Bowes, M. J.; Crossley, A.; Jarvie, H. P.; Jurkschat, K.; Jurgens, M. D.; Lawlor, A. J.; Park, B.; Rowland, P.; Spurgeon, D.; Svendsen, C.; Thompson, I. P.; Barnes, R. J.; Williams, R. J.; Xu, N., An assessment of the fate, behaviour and environmental risk associated with sunscreen TiO₂ nanoparticles in UK field scenarios. *Science of the Total Environment* **2011**, *409*, (13), 2503-2510.
43. Cong, Y.; Banta, G. T.; Selck, H.; Berhanu, D.; Valsami-Jones, E.; Forbes, V. E., Toxic effects and bioaccumulation of nano-, micron- and ionic-Ag in the polychaete, *Nereis diversicolor*. *Aquatic Toxicology* **2011**, *105*, (3-4), 403-411.
44. Al-Salim, N.; Barraclough, E.; Burgess, E.; Clothier, B.; Deurer, M.; Green, S.; Malone, L.; Weir, G., Quantum dot transport in soil, plants, and insects. *Sci. Total Environ.* **2011**, *409*, (17), 3237-48.
45. Hildebrand, H.; Kühnel, D.; Potthoff, A.; Mackenzie, K.; Springer, A.; Schirmer, K., Evaluating the cytotoxicity of palladium/magnetite nano-catalysts intended for wastewater treatment. *Environmental Pollution* **2010**, *158*, (1), 65-73.
46. Gardea-Torresdey, J. L.; Rico, C. M.; White, J. C., Trophic transfer, transformation, and impact of engineered nanomaterials in terrestrial environments. *Environmental Science & Technology* **2014**, *48*, (5), 2526-2540.
47. Musee, N.; Thwala, M.; Nota, N., The antibacterial effects of engineered nanomaterials: implications for wastewater treatment plants. *Journal of Environmental Monitoring* **2011**, *13*, (5), 1164-1183.
48. Nyberg, L.; Turco, R. F.; Nies, L., Assessing the impact of nanomaterials on anaerobic microbial communities. *Environmental Science & Technology* **2008**, *42*, (6), 1938-1943.
49. Costanza, R.; d'Arge, R.; de Groot, R.; Farber, S.; Grasso, M.; Hannon, B.; Limburg, K.; Naeem, S.; O'Neill, R. V.; Paruelo, J.; Raskin, R. G.; Sutton, P.; van den Belt, M., The value of the world's ecosystem services and natural capital. *Nature* **1997**, *387*, (6630), 253-260.
50. Rico, C. M.; Lee, S. C.; Rubenecia, R.; Mukherjee, A.; Hong, J.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Cerium oxide nanoparticles impact yield and modify nutritional parameters in wheat (*Triticum aestivum* L.). *Journal of Agricultural and Food Chemistry* **2014**, *62*, (40), 9669-9675.
51. Rico, C. M.; Barrios, A. C.; Tan, W. J.; Rubenecia, R.; Lee, S. C.; Varela-Ramirez, A.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Physiological and biochemical response of soil-grown barley (*Hordeum vulgare* L.) to cerium oxide nanoparticles. *Environmental Science and Pollution Research* **2015**, *22*, (14), 10551-10558.
52. Kraus, N.; Malmfors, T.; Slovic, P., Intuitive toxicology: expert and lay judgements of chemical risks. *Risk Analysis* **1992**, *12*, (2), 215-232.
53. Holden, P. A.; Klaessig, F.; Turco, R. F.; Priester, J. H.; Rico, C. M.; Avila-Arias, H.; Mortimer, M.; Pacpaco, K.; Gardea-Torresdey, J. L., Evaluation of exposure concentrations used in assessing manufactured nanomaterial environmental hazards: are they relevant? *Environmental Science & Technology* **2014**, *48*, (18), 10541-10551.
54. El Badawy, A. M.; Silva, R. G.; Morris, B.; Scheckel, K. G.; Suidan, M. T.; Tolaymat, T. M., Surface charge-dependent toxicity of silver nanoparticles. *Environmental Science & Technology* **2011**, *45*, (1), 283-287.
55. Corsi, I.; Cherr, G. N.; Lenihan, H. S.; Labille, J.; Hasselov, M.; Canesi, L.; Dondero, F.; Frenzilli, G.; Hristozov, D.; Puentes, V.; Della Torre, C.; Pinsino, A.; Libralato, G.; Marcomini, A.; Sabbioni, E.; Matranga, V., Common strategies and technologies for the ecosafety

assessment and design of nanomaterials entering the marine environment. *ACS Nano* **2014**, 8, (10), 9694-9709.

56. Hendren, C. O.; Lowry, M.; Grieger, K. D.; Money, E. S.; Johnston, J. M.; Wiesner, M. R.; Beaulieu, S. M., Modeling approaches for characterizing and evaluating environmental exposure to engineered nanomaterials in support of risk-based decision making. *Environmental Science & Technology* **2013**, 47, (3), 1190-1205.

57. Handy, R. D.; van den Brink, N.; Chappell, M.; Muhling, M.; Behra, R.; Dusinska, M.; Simpson, P.; Ahtiainen, J.; Jha, A. N.; Seiter, J.; Bednar, A.; Kennedy, A.; Fernandes, T. F.; Riediker, M., Practical considerations for conducting ecotoxicity test methods with manufactured nanomaterials: what have we learnt so far? *Ecotoxicology* **2012**, 21, (4), 933-972.

58. Priester, J. H.; Ge, Y.; Mielke, R. E.; Horst, A. M.; Moritz, S. C.; Espinosa, K.; Gelb, J.; Walker, S. L.; Nisbet, R. M.; An, Y.-J.; Schimel, J. P.; Palmer, R. G.; Hernandez-Viezcás, J. A.; Zhao, L.; Gardea-Torresdey, J. L.; Holden, P. A., Soybean susceptibility to manufactured nanomaterials with evidence for food quality and soil fertility interruption. *Proceedings of the National Academy of Sciences* **2012**, 109, (37), E2451-E2456.

59. Du, W. C.; Gardea-Torresdey, J. L.; Ji, R.; Yin, Y.; Zhu, J. G.; Peralta-Videa, J. R.; Guo, H. Y., Physiological and biochemical changes imposed by CeO₂ nanoparticles on wheat: a life cycle field study. *Environmental Science & Technology* **2015**, 49, (19), 11884-11893.

60. Bour, A.; Mouchet, F.; Silvestre, J.; Gauthier, L.; Pinelli, E., Environmentally relevant approaches to assess nanoparticles ecotoxicity: A review. *Journal of Hazardous Materials* **2015**, 283, 764-777.

61. Murray, K. E.; Thomas, S. M.; Bodour, A. A., Prioritizing research for trace pollutants and emerging contaminants in the freshwater environment. *Environmental Pollution* **2010**, 158, (12), 3462-3471.

62. Petersen, E. J.; Diamond, S. A.; Kennedy, A. J.; Goss, G. G.; Ho, K.; Lead, J.; Hanna, S. K.; Hartmann, N. B.; Hund-Rinke, K.; Mader, B.; Manier, N.; Pandard, P.; Salinas, E. R.; Sayre, P., Adapting OECD aquatic toxicity tests for use with manufactured nanomaterials: key issues and consensus recommendations. *Environmental Science & Technology* **2015**, 49, (16), 9532-9547.

63. Hund-Rinke, K.; Herrchen, M.; Schlich, K. *Integrative test strategy for the environmental assessment of nanomaterials*; (UBA-FB) 002028/E; Environmental Research of the Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety: Dessau-Roßlau, Germany, March, 2015; p 143.

64. Handy, R. D.; Cornelis, G.; Fernandes, T.; Tsyusko, O.; Decho, A.; Sabo-Attwood, T.; Metcalfe, C.; Steevens, J. A.; Klaine, S. J.; Koelmans, A. A.; Horne, N., Ecotoxicity test methods for engineered nanomaterials: practical experiences and recommendations from the bench. *Environmental Toxicology and Chemistry* **2012**, 31, (1), 15-31.

65. OECD, *Preliminary Review of OECD Test Guidelines for Their Applicability to Manufactured Nanomaterials*. OECD Publishing: Paris, 2009; Vol. 15, p 71.

66. OECD, *Ecotoxicology and Environmental Fate of Manufactured Nanomaterials: Test Guidelines*. OECD Publishing: Paris, 2014; Vol. 40, p 84.

67. Kuhnel, D.; Nickel, C., The OECD expert meeting on ecotoxicology and environmental fate - Towards the development of improved OECD guidelines for the testing of nanomaterials. *Science of the Total Environment* **2014**, 472, 347-353.

68. Priester, J. H.; Singhal, A.; Wu, B.; Stucky, G. D.; Holden, P. A., Integrated approach to evaluating the toxicity of novel cysteine-capped silver nanoparticles to *Escherichia coli* and *Pseudomonas aeruginosa*. *Analyst* **2014**, *139*, (5), 954-963.
69. Miller, R. J.; Bennett, S.; Keller, A. A.; Pease, S.; Lenihan, H. S., TiO₂ nanoparticles are phototoxic to marine phytoplankton. *PLoS ONE* **2012**, *7*, (1), e30321.
70. Kaweeteerawat, C.; Chang, C. H.; Roy, K. R.; Liu, R.; Li, R.; Toso, D.; Fischer, H.; Ivask, A.; Ji, Z.; Zink, J. I.; Zhou, Z. H.; Chanfreau, G. F.; Telesca, D.; Cohen, Y.; Holden, P. A.; Nel, A. E.; Godwin, H. A., Cu nanoparticles have different impacts in *Escherichia coli* and *Lactobacillus brevis* than their micro-sized and ionic analogues. *ACS Nano* **2015**, *9*, (7), 7215-7225.
71. Priester, J. H.; Stoimenov, P. K.; Mielke, R. E.; Webb, S. M.; Ehrhardt, C.; Zhang, J. P.; Stucky, G. D.; Holden, P. A., Effects of soluble cadmium salts versus CdSe quantum dots on the growth of planktonic *Pseudomonas aeruginosa*. *Environmental Science & Technology* **2009**, *43*, (7), 2589-2594.
72. Guo, X. K.; Dong, S. P.; Petersen, E. J.; Gao, S. X.; Huang, Q. G.; Mao, L., Biological uptake and depuration of radio-labeled graphene by *Daphnia magna*. *Environmental Science & Technology* **2013**, *47*, (21), 12524-12531.
73. Tervonen, K.; Waissi, G.; Petersen, E. J.; Akkanen, J.; Kukkonen, J. V. K., Analysis of fullerene-C-60 and kinetic measurements for its accumulation and depuration in *Daphnia magna*. *Environmental Toxicology and Chemistry* **2010**, *29*, (5), 1072-1078.
74. Ivask, A.; Suarez, E.; Patel, T.; Boren, D.; Ji, Z. X.; Holden, P.; Telesca, D.; Damoiseaux, R.; Bradley, K. A.; Godwin, H., Genome-wide bacterial toxicity screening uncovers the mechanisms of toxicity of a cationic polystyrene nanomaterial. *Environmental Science & Technology* **2012**, *46*, (4), 2398-2405.
75. Kaweeteerawat, C.; Ivask, A.; Liu, R.; Zhang, H.; Chang, C. H.; Low-Kam, C.; Fischer, H.; Ji, Z.; Pokhrel, S.; Cohen, Y.; Telesca, D.; Zink, J.; Mädler, L.; Holden, P. A.; Nel, A.; Godwin, H., Toxicity of metal oxide nanoparticles in *Escherichia coli* correlates with conduction band and hydration energies. *Environmental Science & Technology* **2015**, *49*, (2), 1105-1112.
76. Chen, C.; Unrine, J. M.; Judy, J. D.; Lewis, R. W.; Guo, J.; McNear, D. H.; Tsyusko, O. V., Toxicogenomic responses of the model legume *Medicago truncatula* to aged biosolids containing a mixture of nanomaterials (TiO₂, Ag, and ZnO) from a pilot wastewater treatment plant. *Environmental Science & Technology* **2015**, *49*, (14), 8759-8768.
77. Gou, N.; Onnis-Hayden, A.; Gu, A. Z., Mechanistic toxicity assessment of nanomaterials by whole-cell-array stress genes expression analysis. *Environmental Science & Technology* **2010**, *44*, (15), 5964-5970.
78. Ma, C.; Chhikara, S.; Xing, B.; Musante, C.; White, J. C.; Dhankher, O. P., Physiological and molecular response of *Arabidopsis thaliana* (L.) to nanoparticle cerium and indium oxide exposure. *ACS Sustainable Chem. Eng.* **2013**, *1*, 768-778.
79. Marmiroli, M.; Pagano, L.; Sardaro, M. L. S.; Villani, M.; Marmiroli, N., Genome-wide approach in *Arabidopsis thaliana* to assess the toxicity of cadmium sulfide quantum dots. *Environmental Science & Technology* **2014**, *48*, (10), 5902-5909.
80. Lan, J.; Gou, N.; Gao, C.; He, M.; Gu, A. Z., Comparative and mechanistic genotoxicity assessment of nanomaterials via a quantitative toxicogenomics approach across multiple species. *Environmental Science & Technology* **2014**, *48*, (21), 12937-12945.
81. Chang, X. J.; Henderson, W. M.; Bouchard, D. C., Multiwalled carbon nanotube dispersion methods affect their aggregation, deposition, and biomarker response. *Environmental Science & Technology* **2015**, *49*, (11), 6645-6653.

82. Wang, Q.; Ebbs, S. D.; Chen, Y. S.; Ma, X. M., Trans-generational impact of cerium oxide nanoparticles on tomato plants. *Metallomics* **2013**, *5*, (6), 753-759.
83. Arndt, D. A.; Chen, J.; Moua, M.; Klaper, R. D., Multigeneration impacts on *Daphnia magna* of carbon nanomaterials with differing core structures and functionalizations. *Environmental Toxicology and Chemistry* **2014**, *33*, (3), 541-547.
84. Petersen, J. E.; Kemp, W. M., Mesocosms: enclosed experimental ecosystems in ocean science. In *Encyclopedia of Ocean Sciences*, second ed.; Steele, J. H.; Turekian, K. K.; Thorpe, S. A., Eds. Academic Press: 2009; pp 732-747.
85. Ge, Y.; Schimel, J. P.; Holden, P. A., Evidence for negative effects of TiO₂ and ZnO nanoparticles on soil bacterial communities. *Environmental Science & Technology* **2011**, *45*, (4), 1659-1664.
86. Pakarinen, K.; Petersen, E. J.; Leppanen, M. T.; Akkanen, J.; Kukkonen, J. V. K., Adverse effects of fullerenes (nC₆₀) spiked to sediments on *Lumbriculus variegatus* (Oligochaeta). *Environmental Pollution* **2011**, *159*, (12), 3750-3756.
87. Ge, Y.; Priester, J. H.; Van De Werfhorst, L. C.; Walker, S. L.; Nisbet, R. M.; An, Y.-J.; Schimel, J. P.; Gardea-Torresdey, J. L.; Holden, P. A., Soybean plants modify metal oxide nanoparticle effects on soil bacterial communities. *Environmental Science & Technology* **2014**, *48*, (22), 13489-13496.
88. Hanna, S. K.; Miller, R. J.; Muller, E. B.; Nisbet, R. M.; Lenihan, H. S., Impact of engineered zinc oxide nanoparticles on the individual performance of *Mytilus galloprovincialis*. *Plos One* **2013**, *8*, (4), 1-7.
89. Hanna, S. K.; Miller, R. J.; Zhou, D. X.; Keller, A. A.; Lenihan, H. S., Accumulation and toxicity of metal oxide nanoparticles in a soft-sediment estuarine amphipod. *Aquatic Toxicology* **2013**, *142*, 441-446.
90. Dai, L.; Syberg, K.; Banta, G. T.; Selck, H.; Forbes, V. E., Effects, uptake, and depuration kinetics of silver oxide and copper oxide nanoparticles in a marine deposit feeder, *Macoma balthica*. *ACS Sustainable Chemistry & Engineering* **2013**, *1*, (7), 760-767.
91. Keller, A. A.; Wang, H. T.; Zhou, D. X.; Lenihan, H. S.; Cherr, G.; Cardinale, B. J.; Miller, R.; Ji, Z. X., Stability and aggregation of metal oxide nanoparticles in natural aqueous matrices. *Environmental Science & Technology* **2010**, *44*, (6), 1962-1967.
92. Echavarri-Bravo, V.; Paterson, L.; Aspray, T. J.; Porter, J. S.; Winson, M. K.; Thornton, B.; Hartl, M. G. J., Shifts in the metabolic function of a benthic estuarine microbial community following a single pulse exposure to silver nanoparticles. *Environmental Pollution* **2015**, *201*, 91-99.
93. Ge, Y.; Schimel, J. P.; Holden, P. A., Identification of soil bacteria susceptible to TiO₂ and ZnO nanoparticles. *Applied & Environmental Microbiology* **2012**, *78*, (18), 6749-6758.
94. Priester, J. H.; Ge, Y.; Chang, V.; Stoimenov, P. K.; Schimel, J. P.; Stucky, G. D.; Holden, P. A., Assessing interactions of hydrophilic nanoscale TiO₂ with soil water. *Journal of Nanoparticle Research* **2013**, *15*, (9), 1-13.
95. Ge, Y.; Priester, J. H.; De Werfhorst, L. C. V.; Schimel, J. P.; Holden, P. A., Potential mechanisms and environmental controls of TiO₂ nanoparticle effects on soil bacterial communities. *Environmental Science & Technology* **2013**, *47*, (24), 14411-14417.
96. Horst, A. M.; Neal, A. C.; Mielke, R. E.; Sislian, P. R.; Suh, W. H.; Mädler, L.; Stucky, G. D.; Holden, P. A., Dispersion of TiO₂ nanoparticle agglomerates by *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* **2010**, *76*, (21), 7292-7298.

97. Rhiem, S.; Riding, M. J.; Baumgartner, W.; Martin, F. L.; Semple, K. T.; Jones, K. C.; Schaffer, A.; Maes, H. M., Interactions of multiwalled carbon nanotubes with algal cells: Quantification of association, visualization of uptake, and measurement of alterations in the composition of cells. *Environmental Pollution* **2015**, *196*, 431-439.
98. Jarvis, T. A.; Miller, R. J.; Lenihan, H. S.; Bielmyer, G. K., Toxicity of ZnO nanoparticles to the copepod *Acartia tonsa*, exposed through a phytoplankton diet. *Environmental Toxicology and Chemistry* **2013**, *32*, (6), 1264-1269.
99. Mielke, R. E.; Priester, J. H.; Werlin, R. A.; Gelb, J.; Horst, A. M.; Orias, E.; Holden, P. A., Differential growth of and nanoscale TiO₂ accumulation in *Tetrahymena thermophila* by direct feeding versus trophic transfer from *Pseudomonas aeruginosa*. *Applied and Environmental Microbiology* **2013**, *79*, (18), 5616-5624.
100. Khan, F. R.; Paul, K. B.; Dybowska, A. D.; Valsami-Jones, E.; Lead, J. R.; Stone, V.; Fernandes, T. F., Accumulation dynamics and acute toxicity of silver nanoparticles to *Daphnia magna* and *Lumbriculus variegatus*: implications for metal modeling approaches. *Environmental Science & Technology* **2015**, *49*, (7), 4389-4397.
101. Kalman, J.; Paul, K. B.; Khan, F. R.; Stone, V.; Fernandes, T. F., Characterisation of bioaccumulation dynamics of three differently coated silver nanoparticles and aqueous silver in a simple freshwater food chain. *Environmental Chemistry* **2015**, *12*, (6), 662-672.
102. Werlin, R.; Priester, J. H.; Mielke, R. E.; Krämer, S.; Jackson, S.; Stoimenov, P. K.; Stucky, G. D.; Cherr, G. N.; Orias, E.; Holden, P. A., Biomagnification of cadmium selenide quantum dots in a simple experimental microbial food chain. *Nature Nanotechnology* **2011**, *6*, (1), 65-71.
103. Yue, Y.; Behra, R.; Sigg, L.; Freire, P. F.; Pillai, S.; Schirmer, K., Toxicity of silver nanoparticles to a fish gill cell line: Role of medium composition. *Nanotoxicology* **2015**, *9*, (1), 54-63.
104. Smulders, S.; Luyts, K.; Brabants, G.; Golanski, L.; Martens, J.; Vanoirbeek, J.; Hoet, P. H. M., Toxicity of nanoparticles embedded in paints compared to pristine nanoparticles, in vitro study. *Toxicology Letters* **2015**, *232*, (2), 333-339.
105. Gottschalk, F.; Nowack, B., The release of engineered nanomaterials to the environment. *Journal of Environmental Monitoring* **2011**, *13*, (5), 1145-1155.
106. Solovitch, N.; Labille, J. r. m.; Rose, J. r. m.; Chaurand, P.; Borschneck, D.; Wiesner, M. R.; Bottero, J.-Y., Concurrent aggregation and deposition of TiO₂ nanoparticles in a sandy porous media. *Environmental Science & Technology* **2010**, *44*, (13), 4897-4902.
107. Auffan, M.; Bottero, J. Y.; Chaneac, C.; Rose, J., Inorganic manufactured nanoparticles: how their physicochemical properties influence their biological effects in aqueous environments. *Nanomedicine* **2010**, *5*, (6), 999-1007.
108. Tong, Z. H.; Bischoff, M.; Nies, L. F.; Myer, P.; Applegate, B.; Turco, R. F., Response of soil microorganisms to as-produced and functionalized single-wall carbon nanotubes (SWNTs). *Environmental Science & Technology* **2012**, *46*, (24), 13471-13479.
109. Simonin, M.; Guyonnet, J. P.; Martins, J. M. F.; Ginot, M.; Richaume, A., Influence of soil properties on the toxicity of TiO₂ nanoparticles on carbon mineralization and bacterial abundance. *Journal of Hazardous Materials* **2015**, *283*, 529-535.
110. Schlich, K.; Klawonn, T.; Terytze, K.; Hund-Rinke, K., Effects of silver nanoparticles and silver nitrate in the earthworm reproduction test. *Environmental Toxicology and Chemistry* **2013**, *32*, (1), 181-188.

111. Lopez-Moreno, M. L.; de la Rosa, G.; Hernandez-Viezcas, J. A.; Castillo-Michel, H.; Botez, C. E.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Evidence of the differential biotransformation and genotoxicity of ZnO and CeO₂ nanoparticles on soybean (*Glycine max*) plants. *Environmental Science & Technology* **2010**, *44*, (19), 7315-7320.
112. Mackay, D.; McCarty, L. S.; Arnot, J. A., Relationships between exposure and dose in aquatic toxicity tests for organic chemicals. *Environmental Toxicology and Chemistry* **2014**, *33*, (9), 2038-2046.
113. Gordon, A. K.; Palmer, C. G., Defining an exposure-response relationship for suspended kaolin clay particulates and aquatic organisms: work toward defining a water quality guideline for suspended solids. *Environmental Toxicology and Chemistry* **2015**, *34*, (4), 907-912.
114. Koelmans, A. A.; Diepens, N. J.; Velzeboer, I.; Besseling, E.; Quik, J. T. K.; van de Meent, D., Guidance for the prognostic risk assessment of nanomaterials in aquatic ecosystems. *Science of The Total Environment* **2015**, *535*, 141-249.
115. Sekine, R.; Brunetti, G.; Donner, E.; Khaksar, M.; Vasilev, K.; Jamting, A. K.; Scheckel, K. G.; Kappen, P.; Zhang, H.; Lombi, E., Speciation and lability of Ag⁺, AgCl⁻, and Ag₂S-nanoparticles in soil determined by X-ray absorption spectroscopy and diffusive gradients in thin films. *Environmental Science & Technology* **2015**, *49*, (2), 897-905.
116. Cui, D.; Zhang, P.; Ma, Y. H.; He, X.; Li, Y. Y.; Zhang, J.; Zhao, Y. C.; Zhang, Z. Y., Effect of cerium oxide nanoparticles on asparagus lettuce cultured in an agar medium. *Environmental Science-Nano* **2014**, *1*, (5), 459-465.
117. Hernandez-Viezcas, J. A.; Castillo-Michel, H.; Andrews, J. C.; Cotte, M.; Rico, C.; Peralta-Videa, J. R.; Ge, Y.; Priester, J. H.; Holden, P. A.; Gardea-Torresdey, J. L., In situ synchrotron X-ray fluorescence mapping and speciation of CeO₂ and ZnO nanoparticles in soil cultivated soybean (*Glycine max*). *ACS Nano* **2013**, *7*, (2), 1415-1423.
118. Zou, X. Y.; Shi, J. P.; Zhang, H. W., Coexistence of silver and titanium dioxide nanoparticles: enhancing or reducing environmental risks? *Aquatic Toxicology* **2014**, *154*, 168-175.
119. Canesi, L.; Frenzilli, G.; Balbi, T.; Bernardeschi, M.; Ciacci, C.; Corsolini, S.; Della Torre, C.; Fabbri, R.; Faleri, C.; Focardi, S.; Guidi, P.; Kocan, A.; Marcomini, A.; Mariottini, M.; Nigro, M.; Pozo-Gallardo, K.; Rocco, L.; Scarcelli, V.; Smerilli, A.; Corsi, I., Interactive effects of n-TiO₂ and 2,3,7,8-TCDD on the marine bivalve *Mytilus galloprovincialis*. *Aquatic Toxicology* **2014**, *153*, 53-65.
120. Della Torre, C.; Balbi, T.; Grassi, G.; Frenzilli, G.; Bernardeschi, M.; Smerilli, A.; Guidi, P.; Canesi, L.; Nigro, M.; Monaci, F.; Scarcelli, V.; Rocco, L.; Focardi, S.; Monopoli, M.; Corsi, I., Titanium dioxide nanoparticles modulate the toxicological response to cadmium in the gills of *Mytilus galloprovincialis*. *Journal of Hazardous Materials* **2015**, *297*, 92-100.
121. Hartmann, N. B.; Von der Kammer, F.; Hofmann, T.; Baalousha, M.; Ottofuelling, S.; Baun, A., Algal testing of titanium dioxide nanoparticles-testing considerations, inhibitory effects and modification of cadmium bioavailability. *Toxicology* **2010**, *269*, (2-3), 190-197.
122. de La Torre-Roche, R.; Hawthorne, J.; Deng, Y. Q.; Xing, B. S.; Cai, W. J.; Newman, L. A.; Wang, C.; Ma, X. M.; White, J. C., Fullerene-enhanced accumulation of p,p'-DDE in agricultural crop species. *Environ. Sci. Technol.* **2012**, *46*, (17), 9315-9323.
123. Hu, X. L.; Liu, J. F.; Mayer, P.; Jiang, G., Impacts of some environmentally relevant parameters on the sorption of polycyclic aromatic hydrocarbons to aqueous suspensions of fullerene. *Environmental Toxicology and Chemistry* **2008**, *27*, (9), 1868-1874.

124. Gai, K.; Shi, B. Y.; Yan, X. M.; Wang, D. S., Effect of dispersion on adsorption of atrazine by aqueous suspensions of fullerenes. *Environmental Science & Technology* **2011**, 45, (14), 5959-5965.
125. Sun, K.; Zhang, Z. Y.; Gao, B.; Wang, Z. Y.; Xu, D. Y.; Jin, J.; Liu, X. T., Adsorption of diuron, fluridone and norflurazon on single-walled and multi-walled carbon nanotubes. *Science of the Total Environment* **2012**, 439, 1-7.
126. Chen, Q. Q.; Yin, D. Q.; Hu, X. L.; Wang, R.; Zhang, C., The effect of nC₆₀ on tissue distribution of ibuprofen in *Cyprinus carpio*. *Science of the Total Environment* **2014**, 496, 453-460.
127. Kelsey, J. W.; White, J. C., Effect of C₆₀ fullerenes on the accumulation of weathered *p,p'*-DDE by plant and earthworm species under single and multispecies conditions. *Environmental Toxicology and Chemistry* **2013**, 32, (5), 1117-1123.
128. Guo, H. Y.; Zhang, Z. Y.; Xing, B. S.; Mukherjee, A.; Musante, C.; White, J. C.; He, L. L., Analysis of silver nanoparticles in antimicrobial products using surface-enhanced Raman spectroscopy (SERS). *Environmental Science & Technology* **2015**, 49, (7), 4317-4324.
129. Zhao, J.; Wang, Z. Y.; Mashayekhi, H.; Mayer, P.; Chefetz, B.; Xing, B. S., Pulmonary surfactant suppressed phenanthrene adsorption on carbon nanotubes through solubilization and competition as examined by passive dosing technique. *Environmental Science & Technology* **2012**, 46, (10), 5369-5377.
130. Wang, F.; Yao, J.; Liu, H. J.; Liu, R. P.; Chen, H. L.; Yi, Z. J.; Yu, Q.; Ma, L.; Xing, B. S., Cu and Cr enhanced the effect of various carbon nanotubes on microbial communities in an aquatic environment. *Journal of Hazardous Materials* **2015**, 292, 137-145.
131. Tian, S. Y.; Zhang, Y. D.; Song, C. Z.; Zhu, X. S.; Xing, B. S., Titanium dioxide nanoparticles as carrier facilitate bioaccumulation of phenanthrene in marine bivalve, ark shell (*Scapharca subcrenata*). *Environmental Pollution* **2014**, 192, 59-64.
132. Tian, S. Y.; Zhang, Y. D.; Song, C. Z.; Zhu, X. S.; Xing, B. S., Bioaccumulation and biotransformation of polybrominated diphenyl ethers in the marine bivalve (*Scapharca subcrenata*): Influence of titanium dioxide nanoparticles. *Marine Pollution Bulletin* **2015**, 90, (1-2), 48-53.
133. Liu, F. F.; Zhao, J.; Wang, S. G.; Du, P.; Xing, B. S., Effects of solution chemistry on adsorption of selected pharmaceuticals and personal care products (PPCPs) by graphenes and carbon nanotubes. *Environmental Science & Technology* **2014**, 48, (22), 13197-13206.
134. De La Torre-Roche, R.; Hawthorne, J.; Deng, Y. Q.; Xing, B. S.; Cai, W. J.; Newman, L. A.; Wang, Q.; Ma, X. M.; Hamdi, H.; White, J. C., Multiwalled carbon nanotubes and C-60 fullerenes differentially impact the accumulation of weathered pesticides in four agricultural plants. *Environmental Science & Technology* **2013**, 47, (21), 12539-12547.
135. Zhang, D.; Pan, B.; Cook, R. L.; Xing, B. S., Multi-walled carbon nanotube dispersion by the adsorbed humic acids with different chemical structures. *Environmental Pollution* **2015**, 196, 292-299.
136. Pan, B.; Zhang, D.; Li, H.; Wu, M.; Wang, Z. Y.; Xing, B. S., Increased adsorption of sulfamethoxazole on suspended carbon nanotubes by dissolved humic acid. *Environmental Science & Technology* **2013**, 47, (14), 7722-7728.
137. Wu, F. C.; Bai, Y. C.; Mu, Y. S.; Pan, B.; Xing, B. S.; Lin, Y., Fluorescence quenching of fulvic acids by fullerene in water. *Environmental Pollution* **2013**, 172, 100-107.
138. Pakarinen, K.; Petersen, E. J.; Alvila, L.; Waissi-Leinonen, G. C.; Akkanen, J.; Leppanen, M. T.; Kukkonen, J. V. K., A screening study on the fate of fullerenes (nC₆₀) and their toxic

1390 implications in natural freshwaters. *Environmental Toxicology and Chemistry* **2013**, 32, (6),
1391 1224-1232.

1392 139. Hüffer, T.; Kah, M.; Hofmann, T.; Schmidt, T. C., How redox conditions and irradiation
1393 affect sorption of PAHs by dispersed fullerenes nC₆₀. *Environmental Science & Technology*
1394 **2013**, 47, (13), 6935-6942.

1395 140. Hou, W. C.; Chowdhury, I.; Goodwin, D. G.; Henderson, W. M.; Fairbrother, D. H.;
1396 Bouchard, D.; Zepp, R. G., Photochemical transformation of graphene oxide in sunlight.
1397 *Environmental Science & Technology* **2015**, 49, (6), 3435-3443.

1398 141. Schlich, K.; Hund-Rinke, K., Influence of soil properties on the effect of silver
1399 nanomaterials on microbial activity in five soils. *Environmental Pollution* **2015**, 196, 321-
1400 330.

1401 142. Adeleye, A. S.; Keller, A. A., Long-term colloidal stability and metal leaching of single
1402 wall carbon nanotubes: Effect of temperature and extracellular polymeric substances.
1403 *Water Research* **2014**, 49, 236-250.

1404 143. Farkas, J.; Peter, H.; Ciesielski, T. M.; Thomas, K. V.; Sommaruga, R.; Salvenmoser, W.;
1405 Weyhenmeyer, G. A.; Tranvik, L. J.; Jenssen, B. M., Impact of TiO₂ nanoparticles on
1406 freshwater bacteria from three Swedish lakes. *Science of The Total Environment* **2015**, 535,
1407 85-93.

1408 144. Sauer, U. G.; Aumann, A.; Ma-Hock, L.; Landsiedel, R.; Wohlleben, W., Influence of
1409 dispersive agent on nanomaterial agglomeration and implications for biological effects in
1410 vivo or in vitro. *Toxicology in Vitro* **2015**, 29, (1), 182-186.

1411 145. Huynh, K. A.; McCaffery, J. M.; Chen, K. L., Heteroaggregation reduces antimicrobial
1412 activity of silver nanoparticles: evidence for nanoparticle-cell proximity effects.
1413 *Environmental Science & Technology Letters* **2014**, 1, (9), 361-366.

1414 146. Navarro, E.; Wagner, B.; Odzak, N.; Sigg, L.; Behra, R., Effects of differently coated
1415 silver nanoparticles on the photosynthesis of *Chlamydomonas reinhardtii*. *Environmental*
1416 *Science & Technology* **2015**, 49, (13), 8041-8047.

1417 147. Juganson, K.; Mortimer, M.; Ivask, A.; Kasemets, K.; Kahru, A., Extracellular
1418 conversion of silver ions into silver nanoparticles by protozoan *Tetrahymena thermophila*.
1419 *Environmental Science-Processes & Impacts* **2013**, 15, (1), 244-250.

1420 148. Ghosh, S.; Pradhan, N. R.; Mashayekhi, H.; Dickert, S.; Thantirige, R.; Tuominen, M. T.;
1421 Tao, S.; Xing, B. S., Binary short-range colloidal assembly of magnetic iron oxides
1422 nanoparticles and fullerene (nC₆₀) in environmental media. *Environmental Science &*
1423 *Technology* **2014**, 48, (20), 12285-12291.

1424 149. Praetorius, A.; Tufenkji, N.; Goss, K.-U.; Scheringer, M.; von der Kammer, F.;
1425 Elimelech, M., The road to nowhere: equilibrium partition coefficients for nanoparticles.
1426 *Environmental Science: Nano* **2014**, 1, (4), 317-323.

1427 150. Zhao, J.; Liu, F. F.; Wang, Z. Y.; Cao, X. S.; Xing, B. S., Heteroaggregation of graphene
1428 oxide with minerals in aqueous phase. *Environmental Science & Technology* **2015**, 49, (5),
1429 2849-2857.

1430 151. Zhang, L. W.; Petersen, E. J.; Zhang, W.; Chen, Y. S.; Cabrera, M.; Huang, Q. G.,
1431 Interactions of C-14-labeled multi-walled carbon nanotubes with soil minerals in water.
1432 *Environmental Pollution* **2012**, 166, 75-81.

1433 152. Petersen, E. J.; Henry, T. B.; Zhao, J.; MacCusprie, R. I.; Kirschling, T. L.; Dobrovolskaia,
1434 M. A.; Hackley, V.; Xing, B.; White, J. C., Identification and avoidance of potential artifacts

and misinterpretations in nanomaterial ecotoxicity measurements. *Environmental Science & Technology* **2014**, *48*, (8), 4226-4246.

153. Stevenson, L. M.; Dickson, H.; Klanjscek, T.; Keller, A. A.; McCauley, E.; Nisbet, R. M., Environmental feedbacks and engineered nanoparticles: mitigation of silver nanoparticle toxicity to *Chlamydomonas reinhardtii* by algal-produced organic compounds. *Plos One* **2013**, *8*, (9), 1-7.

154. Editor, Join the dialogue. *Nature Nanotechnology* **2012**, *7*, (9), 545-545.

155. Atha, D. H.; Wang, H. H.; Petersen, E. J.; Cleveland, D.; Holbrook, R. D.; Jaruga, P.; Dizdaroglu, M.; Xing, B. S.; Nelson, B. C., Copper oxide nanoparticle mediated DNA damage in terrestrial plant models. *Environmental Science & Technology* **2012**, *46*, (3), 1819-1827.

156. Hong, J.; Rico, C. M.; Zhao, L. J.; Adeleye, A. S.; Keller, A. A.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Toxic effects of copper-based nanoparticles or compounds to lettuce (*Lactuca sativa*) and alfalfa (*Medicago sativa*). *Environmental Science-Processes & Impacts* **2015**, *17*, (1), 177-185.

157. Peralta-Videa, J. R.; Hernandez-Viezcas, J. A.; Zhao, L.; Diaz, B. C.; Ge, Y.; Priester, J. H.; Holden, P. A.; Gardea-Torresdey, J. L., Cerium dioxide and zinc oxide nanoparticles alter the nutritional value of soil cultivated soybean plants. *Plant Physiology and Biochemistry* **2014**, *80*, 128-135.

158. Tong, Z.; Bischoff, M.; Nies, L.; Applegate, B.; Turco, R. F., Impact of fullerene (C₆₀) on a soil microbial community. *Environmental Science & Technology* **2007**, *41*, (8), 2985-2991.

159. Garner, K. L.; Suh, S.; Lenihan, H. S.; Keller, A. A., Species sensitivity distributions for engineered nanomaterials. *Environmental Science & Technology* **2015**, *49*, (9), 5753-5759.

160. Juganson, K.; Ivask, A.; Blinova, I.; Mortimer, M.; Kahru, A., NanoE-Tox: New and in-depth database concerning ecotoxicity of nanomaterials. *Beilstein Journal of Nanotechnology* **2015**, *6*, 1788-1804.

161. von der Kammer, F.; Ferguson, P. L.; Holden, P. A.; Masion, A.; Rogers, K. R.; Klaine, S. J.; Koelmans, A. A.; Horne, N.; Unrine, J. M., Analysis of engineered nanomaterials in complex matrices (environment and biota): general considerations and conceptual case studies. *Environmental Toxicology and Chemistry* **2012**, *31*, (1), 32-49.

162. Edgington, A. J.; Petersen, E. J.; Herzing, A. A.; Podila, R.; Rao, A.; Klaine, S. J., Microscopic investigation of single-wall carbon nanotube uptake by *Daphnia magna*. *Nanotoxicology* **2014**, *8*, 2-10.

163. Edgington, A. J.; Roberts, A. P.; Taylor, L. M.; Alloy, M. M.; Reppert, J.; Rao, A. M.; Mao, J. D.; Klaine, S. J., The influence of natural organic matter on the toxicity of multiwalled carbon nanotubes. *Environmental Toxicology and Chemistry* **2010**, *29*, (11), 2511-2518.

164. Servin, A. D.; Morales, M. I.; Castillo-Michel, H.; Hernandez-Viezcas, J. A.; Munoz, B.; Zhao, L. J.; Nunez, J. E.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Synchrotron verification of TiO₂ accumulation in cucumber fruit: a possible pathway of TiO₂ nanoparticle transfer from soil into the food chain. *Environmental Science & Technology* **2013**, *47*, (20), 11592-11598.

165. Zhao, L. J.; Sun, Y. P.; Hernandez-Viezcas, J. A.; Hong, J.; Majumdar, S.; Niu, G. H.; Duarte-Gardea, M.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Monitoring the environmental effects of CeO₂ and ZnO nanoparticles through the life cycle of corn (*Zea mays*) plants and in situ μ -XRF mapping of nutrients in kernels. *Environmental Science & Technology* **2015**, *49*, (5), 2921-2928.

166. Holden, P. A.; Schimel, J. P.; Godwin, H. A., Five reasons to use bacteria when assessing manufactured nanomaterial environmental hazards and fates. *Current Opinion in Biotechnology* **2014**, 27, (0), 73-78.
167. Ong, K. J.; MacCormack, T. J.; Clark, R. J.; Ede, J. D.; Ortega, V. A.; Felix, L. C.; Dang, M. K. M.; Ma, G. B.; Fenniri, H.; Veinot, J. G. C.; Goss, G. G., Widespread nanoparticle-assay interference: implications for nanotoxicity testing. *Plos One* **2014**, 9, (3), 1-9.
168. Horst, A. M.; Vukanti, R.; Priester, J. H.; Holden, P. A., An assessment of fluorescence- and absorbance-based assays to study metal-oxide nanoparticle ROS production and effects on bacterial membranes. *Small* **2013**, 9, (9-10), 1753-1764.
169. Park, S.; Woodhall, J.; Ma, G. B.; Veinot, J. G. C.; Cresser, M. S.; Boxall, A. B. A., Regulatory ecotoxicity testing of engineered nanoparticles: are the results relevant to the natural environment? *Nanotoxicology* **2014**, 8, (5), 583-592.
170. Turco, R. F.; Bischoff, M.; Tong, Z. H.; Nies, L., Environmental implications of nanomaterials: are we studying the right thing? *Current Opinion in Biotechnology* **2011**, 22, (4), 527-532.
171. Schimel, J.; Chadwick, O., What's in a name? The importance of soil taxonomy for ecology and biogeochemistry. *Frontiers in Ecology and the Environment* **2013**, 11, (8), 405-406.
172. Sparks, D. L.; Page, A. L.; Helmke, P. A.; Loeppert, R. H., *Methods of Soil Analysis Part 3—Chemical Methods*. Soil Science Society of America, American Society of Agronomy: Madison, WI, 1996.
173. Ji, Z.; Jin, X.; George, S.; Xia, T.; Meng, H.; Wang, X.; Suarez, E.; Zhang, H.; Hoek, E. M. V.; Godwin, H.; Nel, A. E.; Zink, J. I., Dispersion and stability optimization of TiO₂ nanoparticles in cell culture media. *Environmental Science & Technology* **2010**, 44, (19), 7309-7314.
174. Horst, A. M.; Ji, Z. X.; Holden, P. A., Nanoparticle dispersion in environmentally relevant culture media: a TiO₂ case study and considerations for a general approach. *Journal of Nanoparticle Research* **2012**, 14, (8), 1-14.
175. Maes, H. M.; Stibany, F.; Giefers, S.; Daniels, B.; Deutschmann, B.; Baumgartner, W.; Schaffer, A., Accumulation and distribution of multiwalled carbon nanotubes in zebrafish (*Danio rerio*). *Environmental Science & Technology* **2014**, 48, (20), 12256-12264.
176. Song, L.; Vijver, M. G.; de Snoo, G. R.; Peijnenburg, W., Assessing toxicity of copper nanoparticles across five cladoceran species. *Environmental Toxicology and Chemistry* **2015**, 34, (8), 1863-1869.
177. Tella, M.; Auffan, M.; Brousset, L.; Issartel, J.; Kieffer, I.; Pailles, C.; Morel, E.; Santaella, C.; Angeletti, B.; Artells, E.; Rose, J.; Thiery, A.; Bottero, J. Y., Transfer, transformation, and impacts of ceria nanomaterials in aquatic mesocosms simulating a pond ecosystem. *Environmental Science & Technology* **2014**, 48, (16), 9004-9013.
178. Handy, R. D.; von der Kammer, F.; Lead, J. R.; Hasselov, M.; Owen, R.; Crane, M., The ecotoxicology and chemistry of manufactured nanoparticles. *Ecotoxicology* **2008**, 17, (4), 287-314.
179. Pal, S.; Tak, Y. K.; Song, J. M., Does the antibacterial activity of silver nanoparticles depend on the shape of the nanoparticle? A study of the gram-negative bacterium *Escherichia coli*. *Applied and Environmental Microbiology* **2007**, 73, (6), 1712-1720.

180. Hjorth, R.; Sorensen, S. N.; Olsson, M. E.; Baun, A.; Hartmann, N. B., A certain shade of green: can algal pigments reveal shading effects of nanoparticles? *Integrated Environmental Assessment and Management* **2016**, *12*, (1), 200-202.
181. Rasmussen, K.; González, M.; Kearns, P.; Sintes, J. R.; Rossi, F.; Sayre, P., Review of achievements of the OECD Working Party on Manufactured Nanomaterials' Testing and Assessment Programme. From exploratory testing to test guidelines. *Regulatory Toxicology and Pharmacology* **2016**, *74*, 147-160.
182. Sorensen, S. N.; Hjorth, R.; Delgado, C. G.; Hartmann, N. B.; Baun, A., Nanoparticle ecotoxicity - physical and/or chemical effects? *Integrated Environmental Assessment and Management* **2015**, *11*, (4), 722-724.
183. Calder, A. J.; Dimkpa, C. O.; McLean, J. E.; Britt, D. W.; Johnson, W.; Anderson, A. J., Soil components mitigate the antimicrobial effects of silver nanoparticles towards a beneficial soil bacterium, *Pseudomonas chlororaphis* O6. *Science of the Total Environment* **2012**, *429*, 215-222.
184. Kennedy, A. J.; Chappell, M. A.; Bednar, A. J.; Ryan, A. C.; Laird, J. G.; Stanley, J. K.; Steevens, J. A., Impact of organic carbon on the stability and toxicity of fresh and stored silver nanoparticles. *Environmental Science & Technology* **2012**, *46*, (19), 10772-10780.
185. Zhu, M.; Nie, G.; Meng, H.; Xia, T.; Nel, A.; Zhao, Y., Physicochemical properties determine nanomaterial cellular uptake, transport, and fate. *Accounts of Chemical Research* **2012**, *46*, (3), 622-631.
186. Rodrigues, D. F.; Elimelech, M., Toxic effects of single-walled carbon nanotubes in the development of *E. coli* biofilm. *Environmental Science & Technology* **2010**, *44*, (12), 4583-4589.
187. Saleh, N. B.; Chambers, B.; Aich, N.; Plazas-Tuttle, J.; Phung-Ngoc, H. N.; Kirisits, M. J., Mechanistic lessons learned from studies of planktonic bacteria with metallic nanomaterials: implications for interactions between nanomaterials and biofilm bacteria. *Frontiers in Microbiology* **2015**, *6*, 1-8.
188. Burns, J. M.; Pennington, P. L.; Sisco, P. N.; Frey, R.; Kashiwada, S.; Fulton, M. H.; Scott, G. I.; Decho, A. W.; Murphy, C. J.; Shaw, T. J.; Ferry, J. L., Surface charge controls the fate of Au nanorods in saline estuaries. *Environmental Science & Technology* **2013**, *47*, (22), 12844-12851.
189. Ferry, J. L.; Craig, P.; Hexel, C.; Sisco, P.; Frey, R.; Pennington, P. L.; Fulton, M. H.; Scott, I. G.; Decho, A. W.; Kashiwada, S.; Murphy, C. J.; Shaw, T. J., Transfer of gold nanoparticles from the water column to the estuarine food web. *Nature Nanotechnology* **2009**, *4*, (7), 441-444.
190. Furtado, L. M.; Norman, B. C.; Xenopoulos, M. A.; Frost, P. C.; Metcalfe, C. D.; Hintelmann, H., Environmental fate of silver nanoparticles in boreal lake ecosystems. *Environmental Science & Technology* **2015**, *49*, (14), 8441-8450.
191. Furtado, L. M.; Hoque, M. E.; Mitrano, D. F.; Ranville, J. F.; Cheever, B.; Frost, P. C.; Xenopoulos, M. A.; Hintelmann, H.; Metcalfe, C. D., The persistence and transformation of silver nanoparticles in littoral lake mesocosms monitored using various analytical techniques. *Environmental Chemistry* **2014**, *11*, (4), 419-430.
192. Cleveland, D.; Long, S. E.; Pennington, P. L.; Cooper, E.; Fulton, M. H.; Scott, G. I.; Brewer, T.; Davis, J.; Petersen, E. J.; Wood, L., Pilot estuarine mesocosm study on the environmental fate of silver nanomaterials leached from consumer products. *Science of the Total Environment* **2012**, *421*, 267-272.

193. Lowry, G. V.; Espinasse, B. P.; Badireddy, A. R.; Richardson, C. J.; Reinsch, B. C.; Bryant, L. D.; Bone, A. J.; Deonaraine, A.; Chae, S.; Therezien, M.; Colman, B. P.; Hsu-Kim, H.; Bernhardt, E. S.; Matson, C. W.; Wiesner, M. R., Long-term transformation and fate of manufactured Ag nanoparticles in a simulated large scale freshwater emergent wetland. *Environmental Science & Technology* **2012**, *46*, (13), 7027-7036.
194. Buffet, P. E.; Richard, M.; Caupos, F.; Vergnoux, A.; Perrein-Ettajani, H.; Luna-Acosta, A.; Akcha, F.; Amiard, J. C.; Amiard-Triquet, C.; Guibbolini, M.; Risso-De Faverney, C.; Thomas-Guyon, H.; Reip, P.; Dybowska, A.; Berhanu, D.; Valsami-Jones, E.; Mouneyrac, C., A mesocosm study of fate and effects of CuO nanoparticles on endobenthic species (*Scrobicularia plana*, *Hediste diversicolor*). *Environmental Science & Technology* **2013**, *47*, (3), 1620-1628.
195. Benn, T. M.; Westerhoff, P., Nanoparticle silver released into water from commercially available sock fabrics. *Environmental Science & Technology* **2008**, *42*, (11), 4133-4139.
196. Kiser, M. A.; Westerhoff, P.; Benn, T.; Wang, Y.; Perez-Rivera, J.; Hristovski, K., Titanium nanomaterial removal and release from wastewater treatment plants. *Environmental Science & Technology* **2009**, *43*, (17), 6757-6763.
197. Kiser, M. A.; Ladner, D. A.; Hristovski, K. D.; Westerhoff, P. K., Nanomaterial transformation and association with fresh and freeze-dried wastewater activated sludge: implications for testing protocol and environmental fate. *Environmental Science & Technology* **2012**, *46*, (13), 7046-7053.
198. Kaegi, R.; Voegelin, A.; Ort, C.; Sinnet, B.; Thalmann, B.; Krismer, J.; Hagendorfer, H.; Elumelu, M.; Mueller, E., Fate and transformation of silver nanoparticles in urban wastewater systems. *Water Research* **2013**, *47*, (12), 3866-3877.
199. Kaegi, R.; Voegelin, A.; Sinnet, B.; Zuleeg, S.; Hagendorfer, H.; Burkhardt, M.; Siegrist, H., Behavior of metallic silver nanoparticles in a pilot wastewater treatment plant. *Environmental Science & Technology* **2011**, *45*, (9), 3902-3908.
200. Wang, Y. F.; Westerhoff, P.; Hristovski, K. D., Fate and biological effects of silver, titanium dioxide, and C-60 (fullerene) nanomaterials during simulated wastewater treatment processes. *Journal of Hazardous Materials* **2012**, *201*, 16-22.
201. Westerhoff, P. K.; Kiser, A.; Hristovski, K., Nanomaterial removal and transformation during biological wastewater treatment. *Environmental Engineering Science* **2013**, *30*, (3), 109-117.
202. Westerhoff, P.; Song, G. X.; Hristovski, K.; Kiser, M. A., Occurrence and removal of titanium at full scale wastewater treatment plants: implications for TiO₂ nanomaterials. *Journal of Environmental Monitoring* **2011**, *13*, (5), 1195-1203.
203. Priester, J. H.; Van De Werfhorst, L. C.; Ge, Y.; Adeleye, A. S.; Tomar, S.; Tom, L. M.; Piceno, Y. M.; Andersen, G. L.; Holden, P. A., Effects of TiO₂ and Ag nanoparticles on polyhydroxybutyrate biosynthesis by activated sludge bacteria. *Environmental Science & Technology* **2014**, *48*, (24), 14712-14720.
204. Yang, Y.; Zhang, C. Q.; Hu, Z. Q., Impact of metallic and metal oxide nanoparticles on wastewater treatment and anaerobic digestion. *Environmental Science-Processes & Impacts* **2013**, *15*, (1), 39-48.
205. Benn, T.; Cavanagh, B.; Hristovski, K.; Posner, J. D.; Westerhoff, P., The release of nanosilver from consumer products used in the home. *Journal of Environmental Quality* **2010**, *39*, (6), 1875-1882.

206. Weir, A.; Westerhoff, P.; Fabricius, L.; Hristovski, K.; von Goetz, N., Titanium dioxide nanoparticles in food and personal care products. *Environmental Science & Technology* **2012**, *46*, (4), 2242-2250.
207. Kiser, M. A.; Ryu, H.; Jang, H. Y.; Hristovski, K.; Westerhoff, P., Biosorption of nanoparticles to heterotrophic wastewater biomass. *Water Research* **2010**, *44*, (14), 4105-4114.
208. Judy, J. D.; McNear, D. H.; Chen, C.; Lewis, R. W.; Tsyusko, O. V.; Bertsch, P. M.; Rao, W.; Stegemeier, J.; Lowry, G. V.; McGrath, S. P.; Durenkamp, M.; Unrine, J. M., Nanomaterials in biosolids inhibit nodulation, shift microbial community composition, and result in increased metal uptake relative to bulk/dissolved metals. *Environmental Science & Technology* **2015**, *49*, (14), 8751-8758.
209. Evans, T. D., Biosolids in Europe. In *26th WEF Residuals & Biosolids Conference*, Water Environment Federation: Raleigh, North Carolina, USA, 2012.
210. Barton, L. E.; Auffan, M.; Olivi, L.; Bottero, J. Y.; Wiesner, M. R., Heteroaggregation, transformation and fate of CeO₂ nanoparticles in wastewater treatment. *Environmental Pollution* **2015**, *203*, 122-129.
211. Mosier, A. C.; Li, Z.; Thomas, B. C.; Hettich, R. L.; Pan, C. L.; Banfield, J. F., Elevated temperature alters proteomic responses of individual organisms within a biofilm community. *ISME Journal* **2015**, *9*, (1), 180-194.
212. Speed, D.; Westerhoff, P.; Sierra-Alvarez, R.; Draper, R.; Pantano, P.; Aravamudhan, S.; Chen, K. L.; Hristovski, K.; Herckes, P.; Bi, X.; Yang, Y.; Zeng, C.; Otero-Gonzalez, L.; Mikoryak, C.; Wilson, B. A.; Kosaraju, K.; Tarannum, M.; Crawford, S.; Yi, P.; Liu, X.; Babu, S. V.; Moinpour, M.; Ranville, J.; Montano, M.; Corredor, C.; Posner, J.; Shadman, F., Physical, chemical, and in vitro toxicological characterization of nanoparticles in chemical mechanical planarization suspensions used in the semiconductor industry: towards environmental health and safety assessments. *Environmental Science-Nano* **2015**, *2*, (3), 227-244.
213. Bandyopadhyay, S.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L., Advanced analytical techniques for the measurement of nanomaterials in food and agricultural samples: a review. *Environmental Engineering Science* **2013**, *30*, (3), 118-125.
214. Petersen, E. J.; Zhang, L. W.; Mattison, N. T.; O'Carroll, D. M.; Whelton, A. J.; Uddin, N.; Nguyen, T.; Huang, Q. G.; Henry, T. B.; Holbrook, R. D.; Chen, K. L., Potential release pathways, environmental fate, and ecological risks of carbon nanotubes. *Environmental Science & Technology* **2011**, *45*, (23), 9837-9856.
215. Mader, B. T.; Ellefson, M. E.; Wolf, S. T., Measurements of nanomaterials in environmentally relevant water matrices using liquid nebulization/differential mobility analysis. *Environmental Toxicology and Chemistry* **2015**, *34*, (4), 833-842.
216. Hartmann, G.; Baumgartner, T.; Schuster, M., Influence of particle coating and matrix constituents on the cloud point extraction efficiency of silver nanoparticles (Ag-NPs) and application for monitoring the formation of Ag-NPs from Ag⁺. *Analytical Chemistry* **2014**, *86*, (1), 790-796.
217. Ojeda, C. B.; Rojas, F. S., Separation and preconcentration by cloud point extraction procedures for determination of ions: recent trends and applications. *Microchimica Acta* **2012**, *177*, (1-2), 1-21.
218. McShane, H. V. A.; Sunahara, G. I.; Whalen, J. K.; Hendershot, W. H., Differences in soil solution chemistry between soils amended with nanosized CuO or Cu reference materials:

implications for nanotoxicity tests. *Environmental Science & Technology* **2014**, *48*, (14), 8135-8142.

219. Epel, D.; Luckenbach, T.; Stevenson, C. N.; Macmanus-Spencer, L. A.; Hamdoun, A.; Smital, T., Efflux transporters: newly appreciated roles in protection against pollutants. *Environmental Science & Technology* **2008**, *42*, (11), 3914-3920.

220. Vannuccini, M. L.; Grassi, G.; Leaver, M. J.; Corsi, I., Combination effects of nano-TiO₂ and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on biotransformation gene expression in the liver of European sea bass *Dicentrarchus labrax*. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* **2015**, *176*, 71-78.

221. NRC, *Exposure Science in the 21st Century: A Vision and a Strategy*. The National Academies Press: Washington, DC, 2012; p 196.

222. Lioy, P.; Weisel, C., *Exposure Science: Basic Principles and Applications*. Academic Press: Oxford, 2014; p 1-16.

223. U.S. Environmental Protection Agency, High-throughput exposure forecasting. In Development, O. o. R. a., Ed. U. S. Environmental Protection Agency: 2014; p 2.

224. Blaser, S. A.; Scheringer, M.; MacLeod, M.; Hungerbuhler, K., Estimation of cumulative aquatic exposure and risk due to silver: Contribution of nano-functionalized plastics and textiles. *Science of the Total Environment* **2008**, *390*, (2-3), 396-409.

225. Scheringer, M., Nanoecotoxicology - environmental risks of nanomaterials. *Nature Nanotechnology* **2008**, *3*, (6), 322-323.

226. Johnston, J. M.; Lowry, M.; Beaulieu, S.; Bowles, E., State-of-the-science report on predictive models and modeling approaches for characterizing and evaluating exposure to nanomaterials. In Ecosystems Research Division, N. E. R. L., Ed. U. S. Environmental Protection Agency: Athens, GA, 2010.

227. Lazareva, A.; Keller, A. A., Estimating potential life cycle releases of engineered nanomaterials from wastewater treatment plants. *ACS Sustainable Chemistry & Engineering* **2014**, *2*, (7), 1656-1665.

228. Liu, H. H.; Bilal, M.; Lazareva, A.; Keller, A.; Cohen, Y., Simulation tool for assessing the release and environmental distribution of nanomaterials. *Beilstein Journal of Nanotechnology* **2015**, *6*, 938-951.

229. Meesters, J. A. J.; Koelmans, A. A.; Quik, J. T. K.; Hendriks, A. J.; van de Meent, D., Multimedia modeling of engineered nanoparticles with SimpleBox4nano: model definition and evaluation. *Environmental Science & Technology* **2014**, *48*, (10), 5726-5736.

230. Gottschalk, F.; Lassen, C.; Kjoelholm, J.; Christensen, F.; Nowack, B., Modeling flows and concentrations of nine engineered nanomaterials in the Danish environment. *Int. J. Environ. Res. Public Health* **2015**, *12*, (5), 5581-602.

231. MacLeod, M.; Scheringer, M.; McKone, T. E.; Hungerbuhler, K., The state of multimedia mass-balance modeling in environmental science and decision-making. *Environmental Science & Technology* **2010**, *44*, (22), 8360-8364.

232. Ryan, P. A.; Cohen, Y., Multimedia transport of particle-bound organics - benzo(a)pyrene test case. *Chemosphere* **1986**, *15*, (1), 21-47.

233. Cohen, Y.; Ryan, P. A., Multi-media modeling of environmental transport - trichloroethylene test case. *Environmental Science & Technology* **1985**, *19*, (5), 412-417.

234. Cohen, Y., Organic pollutant transport. *Environmental Science & Technology* **1986**, *20*, (6), 538-544.

235. Utterback, T., Innovation and industrial evolution in manufacturing industries. In *Technology and Global Industry: Companies and Nations in the World Economy*, Guile, B. R.; Brooks, H., Eds. The National Academies Press: Washington, DC, 1987; pp 16-48.
236. Schnoor, J. L.; Sato, C.; McKechnie, D.; Sahoo, D., Processes, Coefficients, and Models for Simulating Toxic Organics and Heavy Metals in Surface Waters. In *Environmental Research Laboratory, O. o. R. a. D., Ed. U.S. Environmental Protection Agency: Athens, GA, 1987; p 303.*
237. Sehgal, A.; Lalatonne, Y.; Berret, J. F.; Morvan, M., Precipitation-redispersion of cerium oxide nanoparticles with poly(acrylic acid): toward stable dispersions. *Langmuir* **2005**, *21*, (20), 9359-9364.
238. Auffan, M.; Masion, A.; Labille, J.; Diot, M. A.; Liu, W.; Olivi, L.; Proux, O.; Ziarelli, F.; Chaurand, P.; Geantet, C.; Bottero, J. Y.; Rose, J., Long-term aging of a CeO₂ based nanocomposite used for wood protection. *Environmental Pollution* **2014**, *188*, 1-7.
239. Siriwardane, I. W. Adsorption of citric acid on cerium oxide nanoparticles (nanoceria) : effects of pH, surface charge and aggregation. University of Iowa, 2012.
240. Mudunkotuwa, I. A.; Grassian, V. H., Citric acid adsorption on TiO₂ nanoparticles in aqueous suspensions at acidic and circumneutral pH: surface coverage, surface speciation, and Its impact on nanoparticle-nanoparticle interactions. *Journal of the American Chemical Society* **2010**, *132*, (42), 14986-14994.
241. Bitter, J. L.; Yang, J.; Beigzadeh Milani, S.; Jafvert, C. T.; Fairbrother, D. H., Transformations of oxidized multiwalled carbon nanotubes exposed to UVC (254 nm) irradiation. *Environmental Science: Nano* **2014**, *1*, (4), 324-337.
242. Hou, W. C.; Jafvert, C. T., Photochemical transformation of aqueous C-60 clusters in sunlight. *Environmental Science & Technology* **2009**, *43*, (2), 362-367.
243. Flores-Cervantes, D. X.; Maes, H. M.; Schäffer, A.; Hollender, J.; Kohler, H.-P. E., Slow biotransformation of carbon nanotubes by horseradish peroxidase. *Environmental Science & Technology* **2014**, *48*, (9), 4826-34.
244. Lu, K.; Huang, Q. G.; Wang, P.; Mao, L., Physicochemical changes of few-layer graphene in peroxidase-catalyzed reactions: characterization and potential ecological effects. *Environmental Science & Technology* **2015**, *49*, (14), 8558-8565.
245. Feng, Y. P.; Lu, K.; Mao, L.; Guo, X. K.; Gao, S. X.; Petersen, E. J., Degradation of ¹⁴C-labeled few layer graphene via Fenton reaction: Reaction rates, characterization of reaction products, and potential ecological effects. *Water Research* **2015**, *84*, 49-57.
246. Zhang, L.; Petersen, E. J.; Habteselassie, M. Y.; Mao, L.; Huang, Q., Degradation of multiwall carbon nanotubes by bacteria. *Environmental Pollution* **2013**, *181*, (0), 335-339.
247. Levard, C. m.; Reinsch, B. C.; Michel, F. M.; Oumahi, C.; Lowry, G. V.; Brown, G. E., Sulfidation processes of PVP-coated silver nanoparticles in aqueous solution: impact on dissolution rate. *Environmental Science & Technology* **2011**, *45*, (12), 5260-5266.
248. Barton, L. E.; Auffan, M.; Bertrand, M.; Barakat, M.; Santaella, C.; Masion, A.; Borschneck, D.; Olivi, L.; Roche, N.; Wiesner, M. R.; Bottero, J. Y., Transformation of pristine and citrate-functionalized CeO₂ nanoparticles in a laboratory-scale activated sludge reactor. *Environmental Science & Technology* **2014**, *48*, (13), 7289-7296.
249. Reed, K.; Cormack, A.; Kulkarni, A.; Mayton, M.; Sayle, D.; Klaessig, F.; Stadler, B., Exploring the properties and applications of nanoceria: is there still plenty of room at the bottom? *Environmental Science-Nano* **2014**, *1*, (5), 390-405.

250. Avanasí, R.; Jackson, W. A.; Sherwin, B.; Mudge, J. F.; Anderson, T. A., C₆₀ fullerene soil sorption, biodegradation, and plant uptake. *Environmental Science & Technology* **2014**, *48*, (5), 2792-2797.
251. Ottofuelling, S.; von der Kammer, F.; Hofmann, T., Commercial titanium dioxide nanoparticles in both natural and synthetic water: comprehensive multidimensional testing and prediction of aggregation behavior. *Environmental Science & Technology* **2011**, *45*, (23), 10045-10052.
252. Nowack, B.; Baalousha, M.; Bornhoft, N.; Chaudhry, Q.; Cornelis, G.; Cotterill, J.; Gondikas, A.; Hasselov, M.; Lead, J.; Mitrano, D. M.; von der Kammer, F.; Wontner-Smith, T., Progress towards the validation of modeled environmental concentrations of engineered nanomaterials by analytical measurements. *Environmental Science-Nano* **2015**, *2*, (5), 421-428.
253. Keller, A. A.; Lazareva, A., Predicted releases of engineered nanomaterials: from global to regional to local. *Environmental Science & Technology Letters* **2014**, *1*, (1), 65-70.
254. Gottschalk, F.; Sun, T. Y.; Nowack, B., Environmental concentrations of engineered nanomaterials: review of modeling and analytical studies. *Environmental Pollution* **2013**, *181*, 287-300.
255. Lee, S.; Bi, X. Y.; Reed, R. B.; Ranville, J. F.; Herckes, P.; Westerhoff, P., Nanoparticle size detection limits by single particle ICP-MS for 40 elements. *Environmental Science & Technology* **2014**, *48*, (17), 10291-10300.
256. Montano, M. D.; Lowry, G. V.; von der Kammer, F.; Blue, J.; Ranville, J. F., Current status and future direction for examining engineered nanoparticles in natural systems. *Environmental Chemistry* **2014**, *11*, (4), 351-366.
257. Bustos, A. R. M.; Petersen, E. J.; Possolo, A.; Winchester, M. R., Post hoc interlaboratory comparison of single particle ICP-MS size measurements of NIST gold nanoparticle reference materials. *Analytical Chemistry* **2015**, *87*, (17), 8809-8817.
258. Sanchís, J.; Božović, D.; Al-Harbi, N. A.; Silva, L. F.; Farré, M.; Barceló, D., Quantitative trace analysis of fullerenes in river sediment from Spain and soils from Saudi Arabia. *Analytical and Bioanalytical Chemistry* **2013**, *405*, (18), 5915-5923.
259. Sanchís, J.; Berrojalbiz, N.; Caballero, G.; Dachs, J.; Farré, M.; Barceló, D., Occurrence of aerosol-bound fullerenes in the Mediterranean sea atmosphere. *Environmental Science & Technology* **2012**, *46*, (3), 1335-1343.
260. Farré, M.; Pérez, S.; Gajda-Schrantz, K.; Osorio, V.; Kantiani, L.; Ginebreda, A.; Barceló, D., First determination of C₆₀ and C₇₀ fullerenes and N-methylfulleropyrrolidine C₆₀ on the suspended material of wastewater effluents by liquid chromatography hybrid quadrupole linear ion trap tandem mass spectrometry. *Journal of Hydrology* **2010**, *383*, (1-2), 44-51.
261. Sanchís, J.; Oliveira, L. F. S.; de Leão, F. B.; Farré, M.; Barceló, D., Liquid chromatography-atmospheric pressure photoionization-Orbitrap analysis of fullerene aggregates on surface soils and river sediments from Santa Catarina (Brazil). *Science of the Total Environment* **2015**, *505*, 172-179.
262. Luo, Z. X.; Wang, Z. H.; Li, Q. Z.; Pan, Q. K.; Yan, C. Z.; Liu, F., Spatial distribution, electron microscopy analysis of titanium and its correlation to heavy metals: occurrence and sources of titanium nanomaterials in surface sediments from Xiamen Bay, China. *Journal of Environmental Monitoring* **2011**, *13*, (4), 1046-1052.
263. Barton, L. E.; Therezien, M.; Auffan, M.; Bottero, J. Y.; Wiesner, M. R., Theory and methodology for determining nanoparticle affinity for heteroaggregation in environmental

matrices using batch measurements. *Environmental Engineering Science* **2014**, *31*, (7), 421-427.

264. Praetorius, A.; Labille, J.; Scheringer, M.; Thill, A.; Hungerbuhler, K.; Bottero, J. Y., Heteroaggregation of titanium dioxide nanoparticles with model natural colloids under environmentally relevant conditions. *Environmental Science & Technology* **2014**, *48*, (18), 10690-10698.

265. Bouchard, D.; Chang, X.; Chowdhury, I., Heteroaggregation of multiwalled carbon nanotubes with sediments. *Environmental Nanotechnology, Monitoring & Management* **2015**, *4*, 42-50.

266. Sabo-Attwood, T.; Unrine, J. M.; Stone, J. W.; Murphy, C. J.; Ghoshroy, S.; Blom, D.; Bertsch, P. M.; Newman, L. A., Uptake, distribution and toxicity of gold nanoparticles in tobacco (*Nicotiana xanthi*) seedlings. *Nanotoxicology* **2012**, *6*, 353-360.

267. Novak, P. J.; Arnold, W. A.; Blazer, V. S.; Halden, R. U.; Klaper, R. D.; Kolpin, D. W.; Kriebel, D.; Love, N. G.; Martinovic-Weigelt, D.; Patisaul, H. B.; Snyder, S. A.; vom Saal, F. S.; Weisbrod, A. V.; Swackhamer, D. L., On the need for a national (US) research program to elucidate the potential risks to human health and the environment posed by contaminants of emerging concern. *Environmental Science & Technology* **2011**, *45*, (9), 3829-3830.

268. Sturla, S. J.; Boobis, A. R.; FitzGerald, R. E.; Hoeng, J.; Kavlock, R. J.; Schirmer, K.; Whelan, M.; Wilks, M. F.; Peitsch, M. C., Systems toxicology: from basic research to risk assessment. *Chemical Research in Toxicology* **2014**, *27*, (3), 314-329.

269. Nel, A.; Xia, T.; Meng, H.; Wang, X.; Lin, S. J.; Ji, Z. X.; Zhang, H. Y., Nanomaterial toxicity testing in the 21st century: use of a predictive toxicological approach and high-throughput screening. *Accounts of Chemical Research* **2013**, *46*, (3), 607-621.

270. Kashiwada, S., Distribution of nanoparticles in the see-through medaka (*Oryzias latipes*). *Environmental Health Perspectives* **2006**, *114*, (11), 1697-1702.

271. Doudrick, K.; Corson, N.; Oberdorster, G.; Eder, A. C.; Herckes, P.; Halden, R. U.; Westerhoff, P., Extraction and quantification of carbon nanotubes in biological matrices with application to rat lung tissue. *ACS Nano* **2013**, *7*, (10), 8849-8856.

272. Arndt, D. A.; Moua, M.; Chen, J.; Klaper, R. D., Core structure and surface functionalization of carbon nanomaterials alter impacts to daphnid mortality, reproduction, and growth: acute assays do not predict chronic exposure impacts. *Environmental Science & Technology* **2013**, *47*, (16), 9444-9452.

273. Ivask, A.; ElBadawy, A.; Kaweeteerawat, C.; Boren, D.; Fischer, H.; Ji, Z. X.; Chang, C. H.; Liu, R.; Tolaymat, T.; Telesca, D.; Zink, J. I.; Cohen, Y.; Holden, P. A.; Godwin, H. A., Toxicity mechanisms in *Escherichia coli* vary for silver nanoparticles and differ from ionic silver. *ACS Nano* **2014**, *8*, (1), 374-386.

274. Maurer-Jones, M. A.; Gunsolus, I. L.; Meyer, B. M.; Christenson, C. J.; Haynes, C. L., Impact of TiO₂ nanoparticles on growth, biofilm formation, and flavin secretion in *Shewanella oneidensis*. *Analytical Chemistry* **2013**, *85*, (12), 5810-5818.

275. OECD, *Guidance Document on Aquatic Toxicity Testing of Difficult Substances and Mixtures*. OECD Publishing: Paris, 2002; p 53.

276. Boncel, S.; Kyziol-Komosinska, J.; Krzyewska, I.; Czupiol, J., Interactions of carbon nanotubes with aqueous/aquatic media containing organic/inorganic contaminants and selected organisms of aquatic ecosystems - A review. *Chemosphere* **2015**, *136*, 211-221.

277. Auffan, M.; Bertin, D.; Chaurand, P.; Pailles, C.; Dominici, C.; Rose, J.; Bottero, J.-Y.; Thiery, A., Role of molting on the biodistribution of CeO₂ nanoparticles within *Daphnia pulex*. *Water Research* **2013**, *47*, (12), 3921-3930.
278. Long, Z.; Ji, J.; Yang, K.; Lin, D.; Wu, F., Systematic and quantitative investigation of the mechanism of carbon nanotubes' toxicity toward algae. *Environmental Science & Technology* **2012**, *46*, (15), 8458-8466.
279. Zhang, H.; Ji, Z.; Xia, T.; Meng, H.; Low-Kam, C.; Liu, R.; Pokhrel, S.; Lin, S.; Wang, X.; Liao, Y.-P.; Wang, M.; Li, L.; Rallo, R.; Damoiseaux, R.; Telesca, D.; Mädler, L.; Cohen, Y.; Zink, J. I.; Nel, A. E., Use of metal oxide nanoparticle band gap to develop a predictive paradigm for oxidative stress and acute pulmonary inflammation. *ACS Nano* **2012**, *6*, (5), 4349-4368.
280. U.S. Environmental Protection Agency Ecological Risk Assessment. <http://www.epa.gov/risk/ecological-risk.htm> (September 6, 2015),
281. Ankley, G. T.; Bennett, R. S.; Erickson, R. J.; Hoff, D. J.; Hornung, M. W.; Johnson, R. D.; Mount, D. R.; Nichols, J. W.; Russom, C. L.; Schmieder, P. K.; Serrano, J. A.; Tietge, J. E.; Villeneuve, D. L., Adverse outcome pathways: a conceptual framework to support ecotoxicology research and risk assessment. *Environmental Toxicology and Chemistry* **2010**, *29*, (3), 730-741.
282. Kramer, V. J.; Etterson, M. A.; Hecker, M.; Murphy, C. A.; Roesijadi, G.; Spade, D. J.; Spromberg, J. A.; Wang, M.; Ankley, G. T., Adverse outcome pathways and ecological risk assessment bridging to population-level effects. *Environmental Toxicology and Chemistry* **2011**, *30*, (1), 64-76.
283. Jager, T.; Zimmer, E. I., Simplified dynamic energy budget model for analysing ecotoxicity data. *Ecological Modelling* **2012**, *225*, 74-81.
284. Klanjscek, T.; Nisbet, R. M.; Priester, J. H.; Holden, P. A., Modeling physiological processes that relate toxicant exposure and bacterial population dynamics. *PLoS ONE* **2012**, *7*, (2), 1-12.
285. Klanjscek, T.; Nisbet, R. M.; Priester, J. H.; Holden, P. A., Dynamic energy budget approach to modeling mechanisms of CdSe quantum dot toxicity. *Ecotoxicology* **2013**, *22*, (2), 319-330.
286. Schierz, A.; Parks, A. N.; Washburn, K. M.; Chandler, G. T.; Ferguson, P. L., Characterization and quantitative analysis of single-walled carbon nanotubes in the aquatic environment using near-infrared fluorescence Spectroscopy. *Environmental Science & Technology* **2012**, *46*, (22), 12262-12271.
287. Bisesi, J. H.; Merten, J.; Liu, K.; Parks, A. N.; Afrooz, A.; Glenn, J. B.; Klaine, S. J.; Kane, A. S.; Saleh, N. B.; Ferguson, P. L.; Sabo-Attwood, T., Tracking and quantification of single-walled carbon nanotubes in fish using near infrared fluorescence. *Environmental Science & Technology* **2014**, *48*, (3), 1973-1983.
288. Schierz, A.; Espinasse, B.; Wiesner, M. R.; Bisesi, J. H.; Sabo-Attwood, T.; Ferguson, P. L., Fate of single walled carbon nanotubes in wetland ecosystems. *Environmental Science-Nano* **2014**, *1*, (6), 574-583.
289. Parks, A. N.; Chandler, G. T.; Ho, K. T.; Burgess, R. M.; Ferguson, P. L., Environmental biodegradability of ¹⁴C single-walled carbon nanotubes by *Trametes vericolor* and natural microbial cultures found in New Bedford Harbor sediment and aerated wastewater treatment plant sludge. *Environmental Toxicology and Chemistry* **2015**, *34*, (2), 247-251.
290. Parks, A. N.; Portis, L. M.; Schierz, P. A.; Washburn, K. M.; Perron, M. M.; Burgess, R. M.; Ho, K. T.; Chandler, G. T.; Ferguson, P. L., Bioaccumulation and toxicity of single-walled

carbon nanotubes to benthic organisms at the base of the marine food chain. *Environmental Toxicology and Chemistry* **2013**, 32, (6), 1270-1277.

291. Parks, A. N.; Burgess, R. M.; Ho, K. T.; Ferguson, P. L., On the likelihood of single-walled carbon nanotubes causing adverse marine ecological effects. *Integrated Environmental Assessment and Management* **2014**, 10, (3), 472-474.

292. Ferguson, P. L.; Chandler, G. T.; Templeton, R. C.; Demarco, A.; Scrivens, W. A.; Englehart, B. A., Influence of sediment-amendment with single-walled carbon nanotubes and diesel soot on bioaccumulation of hydrophobic organic contaminants by benthic invertebrates. *Environmental Science & Technology* **2008**, 42, (10), 3879-3885.

293. Parks, A. N.; Chandler, G. T.; Portis, L. M.; Sullivan, J. C.; Perron, M. M.; Cantwell, M. G.; Burgess, R. M.; Ho, K. T.; Ferguson, P. L., Effects of single-walled carbon nanotubes on the bioavailability of PCBs in field-contaminated sediments. *Nanotoxicology* **2014**, 8, 111-117.

294. Warheit, D. B.; Donner, E. M., How meaningful are risk determinations in the absence of a complete dataset? Making the case for publishing standardized test guideline and 'no effect' studies for evaluating the safety of nanoparticulates versus spurious 'high effect' results from single investigative studies. *Science and Technology of Advanced Materials* **2015**, 16, (3), 1-5.

295. Oomen, A. G.; Bleeker, E. A. J.; Bos, P. M. J.; van Broekhuizen, F.; Gottardo, S.; Groenewold, M.; Hristozov, D.; Hund-Rinke, K.; Irfan, M. A.; Marcomini, A.; Peijnenburg, W.; Rasmussen, K.; Jimenez, A. S.; Scott-Fordsmand, J. J.; van Tongeren, M.; Wiench, K.; Wohlleben, W.; Landsiedel, R., Grouping and read-across approaches for risk assessment of nanomaterials. *Int. J. Environ. Res. Public Health* **2015**, 12, (10), 13415-13434.

296. Walser, T.; Studer, C., Sameness: The regulatory crux with nanomaterial identity and grouping schemes for hazard assessment. *Regulatory Toxicology and Pharmacology* **2015**, 72, (3), 569-571.

297. OECD, Developments in Delegations on the Safety of Manufactured Nanomaterials - Tour De Table. In *Series on the Safety of Manufactured Nanomaterials*, OECD: Paris, 2015.

298. Hristozov, D. R.; Zabeo, A.; Foran, C.; Isigonis, P.; Critto, A.; Marcomini, A.; Linkov, I., A weight of evidence approach for hazard screening of engineered nanomaterials. *Nanotoxicology* **2014**, 8, (1), 72-87.

299. Environment Canada, Screening Assessment for the Challenge: Carbon Black. In Canada, H., Ed. Government of Canada: 2013; p 88.

300. Nel, A. E.; Parak, W. J.; Chan, W. C. W.; Xia, T.; Hersam, M. C.; Brinker, C. J.; Zink, J. I.; Pinkerton, K. E.; Baer, D. R.; Weiss, P. S., Where are we heading in nanotechnology environmental health and safety and materials characterization? *ACS Nano* **2015**, 9, (6), 5627-5630.

1926

Table 1: Summary recommendations regarding exposure conditions used in assessing ENM hazard potentials for model organisms	
	Develop logically conceived, plausible exposure and receptor scenarios in designing experiments, addressing background chemical, physical, and biotic conditions, biological endpoints, and exposure periods.
	Thoroughly characterize exposure media, including natural soils, waters, and sediments, and provide metadata to allow cross-comparisons and for mining influential factors across studies.
	Select ENM forms—including ENM mixtures or other co-contaminants—and scale exposure concentrations to logically well-defined exposure scenarios, or to scenario-independent mechanistic and process-based modeling goals, or to best available analytical instrumentation limitations.
	To the fullest extent possible, characterize ENMs under the exposure conditions and over the time frame of hazard assessments to allow homogeneous exposure or understanding bioavailability and relating endpoint measures to effective forms.
	Examine and choose multiple endpoints for ENM physicochemical characterization, toxicity quantification, and toxicant characterization, subject to the exposure scenario or similar context.
	Quantify body burden and determine compartmentalization of ENMs and transformation products in receptors, to allow comparing effective doses to measured biological responses. Carefully consider body burden assay approaches to avoid artifacts.
	Adopt appropriate experimental control treatments for media additives such as dispersants, and for ENM transformation products that are expected during hazard assessment.
	Incorporate appropriate rapid screening approaches, as prioritized tiers in an ecological hazard assessment hierarchy (Fig. 1).
	Adapt these approaches in response to relevant knowledge generation.

1927


1928

1929


Table 2: Summary recommendations regarding exposure and design considerations in mesocosm assessment of ENM environmental hazards	
	Establish mesocosms only to address clearly defined questions and hypotheses, including those motivated by testing results from lower complexity microcosms, and as related to expected exposure scenarios.
	Design and operate mesocosms using established authoritative principles, and aim to minimize artifacts.
	Follow the recommendations for assessing hazard potentials in using individualized or microcosm experiments (Table 1)
	Make experiment and exposure durations sufficiently long to best represent the exposure scenario underpinning the mesocosm design.
	Anticipate and address challenges in detecting and quantifying ENMs in complex media and myriad receptors in mesocosms, such that mass balances can be performed, and bioaccumulation and trophic transfer quantified.
	Attend to issues of ENMs aging, or otherwise significantly transforming, over the duration of mesocosm operation, and control for, or otherwise assess, specific outcomes related to aged materials.
	Consider assessing effects and ENM interactions with co-contaminants, since they are realistic constituents of most compartments into which ENMs are released.
	Anticipate, recognize, and plan to assess secondary effects on non-target receptors that contribute to ecological processes in complex systems represented by mesocosms.
	Increasingly use sensitive, e.g., omics type, endpoints as they are responsive to questions and hypotheses motivating mesocosm studies.

1930
1931
1932
1933

1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956



1957



1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971


1972

1973

1974

Figure 2: Interactive “tiers” of ENM ecological hazard assessment. Screening studies (left) are recommended starting points for ENM assessments, and are typically conducted under laboratory conditions, using microcosms, batch reactors, or microplates, depending on the experiment. Examples include assessing organismal to population growth effects, bioavailability (including to communities), ENM physical behavior, and trophic transfer. If screening assay results warrant, mesocosms (upper right box) may be initiated to simulate actual environmental conditions in longer-term experiments, to determine the potential for ENMs to impinge on ecosystems. Screening assay results may also motivate determining mechanisms (lower right box) of observed effects on cells or macromolecules and to characterize biochemical, physical, and chemical interactions of ENMs with biological receptors. Knowledge gained within each tier is used to refine the approaches in the other tiers, thereby improving the relevance of each activity. Results inform development of dynamic process-based mathematical models (curved lines linking across tiers) of biological effects.

1975
1976
1977




1978
1979

Figure 3: Conceptual exposure and effects assessments in ENM environmental risk assessment. Exposure derives from far-field emissions, transport, and transformation processes leading to environmental accumulation (bottom “wedge” of ENM complexity and concentration) of either more complex mixtures, or of specific ENMs or transformation products. Top “wedges” depict that either ENM complexity or ENM concentration can increase or decrease along the path of far-field ENM transport. At biological receptors, adverse effects are predicated on near-field exposures. Homo- and heteroaggregation (of multiple particles, here depicted as two) are particle-specific phenomena that may prevent near-field exposure. Biotic responses can influence bioavailability, and thus near-field exposures. Direct effects to biota may manifest across all levels of biological organization (subcellular, to individual, population, community, and ecosystem); effects can also be indirect, e.g., from physical effects of ENMs on nutrient availability.

1989 **TOC Art (for Table of Contents Only)**

1990

1991



1992

1993

1994

1995

1996